

Fractions



Arithmetic of Fractions

Multiplication

It is very easy to multiply two fractions together: simply multiply together the numerators and denominators:

Arithmetic of Fractions

Multiplication

It is very easy to multiply two fractions together: simply multiply together the numerators and denominators:

$$\left(\frac{a}{b}\right) \times \left(\frac{c}{d}\right) = \frac{a \times c}{b \times d} = \frac{ac}{bd}$$

or, using the dot notation for multiplication:

Arithmetic of Fractions

Multiplication

It is very easy to multiply two fractions together: simply multiply together the numerators and denominators:

$$\left(\frac{a}{b}\right) \times \left(\frac{c}{d}\right) = \frac{a \times c}{b \times d} = \frac{ac}{bd}$$

or, using the dot notation for multiplication:

$$\left(\frac{a}{b}\right) \cdot \left(\frac{c}{d}\right) = \frac{a \cdot c}{b \cdot d} = \frac{ac}{bd}$$

Example 1:

$$\left(\frac{5}{8}\right) \times \left(\frac{7}{3}\right) =$$

Example 1:

$$\left(\frac{5}{8}\right) \times \left(\frac{7}{3}\right) = \frac{5 \times 7}{8 \times 3} =$$

Example 1:

$$\left(\frac{5}{8}\right) \times \left(\frac{7}{3}\right) = \frac{5 \times 7}{8 \times 3} = \frac{35}{24}$$

Example 1:

$$\left(\frac{5}{8}\right) \times \left(\frac{7}{3}\right) = \frac{5 \times 7}{8 \times 3} = \frac{35}{24}$$

Suppose $c = x$ and $d = 1$, so that $\frac{c}{d} = \frac{x}{1} = x$. Then

Example 1:

$$\left(\frac{5}{8}\right) \times \left(\frac{7}{3}\right) = \frac{5 \times 7}{8 \times 3} = \frac{35}{24}$$

Suppose $c = x$ and $d = 1$, so that $\frac{c}{d} = \frac{x}{1} = x$. Then

$$\left(\frac{a}{b}\right) \cdot x =$$

Example 1:

$$\left(\frac{5}{8}\right) \times \left(\frac{7}{3}\right) = \frac{5 \times 7}{8 \times 3} = \frac{35}{24}$$

Suppose $c = x$ and $d = 1$, so that $\frac{c}{d} = \frac{x}{1} = x$. Then

$$\left(\frac{a}{b}\right) \cdot x = \left(\frac{a}{b}\right) \cdot \left(\frac{x}{1}\right) =$$

Example 1:

$$\left(\frac{5}{8}\right) \times \left(\frac{7}{3}\right) = \frac{5 \times 7}{8 \times 3} = \frac{35}{24}$$

Suppose $c = x$ and $d = 1$, so that $\frac{c}{d} = \frac{x}{1} = x$. Then

$$\left(\frac{a}{b}\right) \cdot x = \left(\frac{a}{b}\right) \cdot \left(\frac{x}{1}\right) = \frac{ax}{b(1)} =$$

Example 1:

$$\left(\frac{5}{8}\right) \times \left(\frac{7}{3}\right) = \frac{5 \times 7}{8 \times 3} = \frac{35}{24}$$

Suppose $c = x$ and $d = 1$, so that $\frac{c}{d} = \frac{x}{1} = x$. Then

$$\left(\frac{a}{b}\right) \cdot x = \left(\frac{a}{b}\right) \cdot \left(\frac{x}{1}\right) = \frac{ax}{b(1)} = \frac{ax}{b},$$

so multiplying a fraction by x has the same effect as multiplying the numerator of the fraction by x .

Example 2:

$$\frac{2}{3}\pi =$$

Example 1:

$$\left(\frac{5}{8}\right) \times \left(\frac{7}{3}\right) = \frac{5 \times 7}{8 \times 3} = \frac{35}{24}$$

Suppose $c = x$ and $d = 1$, so that $\frac{c}{d} = \frac{x}{1} = x$. Then

$$\left(\frac{a}{b}\right) \cdot x = \left(\frac{a}{b}\right) \cdot \left(\frac{x}{1}\right) = \frac{ax}{b(1)} = \frac{ax}{b},$$

so multiplying a fraction by x has the same effect as multiplying the numerator of the fraction by x .

Example 2:

$$\frac{2}{3}\pi = \frac{2\pi}{3}$$

Example 3: Combine $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right)$ into one fraction.

Example 3: Combine $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right)$ into one fraction.

Solution: $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right) =$

Example 3: Combine $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right)$ into one fraction.

Solution: $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right) = \frac{(7)(-3)}{5(8)} =$

Example 3: Combine $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right)$ into one fraction.

Solution: $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right) = \frac{(7)(-3)}{5(8)} = -\frac{21}{40}$

Example 4: Combine $\left(\frac{10}{3}\right) \cdot \left(\frac{9}{4}\right)$ into one fraction.

Example 3: Combine $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right)$ into one fraction.

Solution: $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right) = \frac{(7)(-3)}{5(8)} = -\frac{21}{40}$

Example 4: Combine $\left(\frac{10}{3}\right) \cdot \left(\frac{9}{4}\right)$ into one fraction.

Solution: $\left(\frac{10}{3}\right) \cdot \left(\frac{9}{4}\right)$ =(factoring 10, 9, and 4, and cancelling)

Example 3: Combine $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right)$ into one fraction.

Solution: $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right) = \frac{(7)(-3)}{5(8)} = -\frac{21}{40}$

Example 4: Combine $\left(\frac{10}{3}\right) \cdot \left(\frac{9}{4}\right)$ into one fraction.

Solution: $\left(\frac{10}{3}\right) \cdot \left(\frac{9}{4}\right)$ =(factoring 10, 9, and 4, and
cancelling)

$$\frac{(2 \times 5)(3 \times 3)}{3(2 \times 2)} =$$

Example 3: Combine $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right)$ into one fraction.

Solution: $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right) = \frac{(7)(-3)}{5(8)} = -\frac{21}{40}$

Example 4: Combine $\left(\frac{10}{3}\right) \cdot \left(\frac{9}{4}\right)$ into one fraction.

Solution: $\left(\frac{10}{3}\right) \cdot \left(\frac{9}{4}\right)$ =(factoring 10, 9, and 4, and cancelling)

$$\frac{(2 \times 5)(3 \times 3)}{3(2 \times 2)} = \frac{(5)(3)}{2} =$$

Example 3: Combine $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right)$ into one fraction.

Solution: $\left(\frac{7}{5}\right) \cdot \left(\frac{-3}{8}\right) = \frac{(7)(-3)}{5(8)} = -\frac{21}{40}$

Example 4: Combine $\left(\frac{10}{3}\right) \cdot \left(\frac{9}{4}\right)$ into one fraction.

Solution: $\left(\frac{10}{3}\right) \cdot \left(\frac{9}{4}\right)$ =(factoring 10, 9, and 4, and cancelling)

$$\frac{(2 \times 5)(3 \times 3)]}{3(2 \times 2)} = \frac{(5)(3)}{2} = \frac{15}{2}$$

Addition

Case 1: Same Denominator:

To add two fractions with the same denominator, *add the numerators together*, i.e.,

$$\frac{a}{h} + \frac{b}{h} = \frac{a+b}{h}$$

(Consider: two-sixths of a pizza plus one-sixth of a pizza equals three-sixths, or one-half of a pizza.)

Case 2: Different Denominators:

For fractions with different denominators, say $\frac{a}{b}$ and $\frac{c}{d}$, we must first alter the fractions to have a **common denominator**, say $h = bd$. We do this by multiplications by 1, where we write 1 as an appropriate fraction:

$$\frac{a}{b} + \frac{c}{d} =$$

Case 2: Different Denominators:

For fractions with different denominators, say $\frac{a}{b}$ and $\frac{c}{d}$, we must first alter the fractions to have a **common denominator**, say $h = bd$. We do this by multiplications by 1, where we write 1 as an appropriate fraction:

$$\frac{a}{b} + \frac{c}{d} = \frac{a}{b} \cdot 1 + 1 \cdot \frac{c}{d} =$$

Case 2: Different Denominators:

For fractions with different denominators, say $\frac{a}{b}$ and $\frac{c}{d}$, we must first alter the fractions to have a **common denominator**, say $h = bd$. We do this by multiplications by 1, where we write 1 as an appropriate fraction:

$$\frac{a}{b} + \frac{c}{d} = \frac{a}{b} \cdot 1 + 1 \cdot \frac{c}{d} = \left(\frac{a}{b}\right) \cdot \left(\frac{d}{d}\right) + \left(\frac{b}{b}\right) \cdot \left(\frac{c}{d}\right) =$$

Case 2: Different Denominators:

For fractions with different denominators, say $\frac{a}{b}$ and $\frac{c}{d}$, we must first alter the fractions to have a **common denominator**, say $h = bd$. We do this by multiplications by 1, where we write 1 as an appropriate fraction:

$$\frac{a}{b} + \frac{c}{d} = \frac{a}{b} \cdot 1 + 1 \cdot \frac{c}{d} = \left(\frac{a}{b}\right) \cdot \left(\frac{d}{d}\right) + \left(\frac{b}{b}\right) \cdot \left(\frac{c}{d}\right) = \frac{ad}{bd} + \frac{bc}{bd} =$$

Case 2: Different Denominators:

For fractions with different denominators, say $\frac{a}{b}$ and $\frac{c}{d}$, we must first alter the fractions to have a **common denominator**, say $h = bd$. We do this by multiplications by 1, where we write 1 as an appropriate fraction:

$$\frac{a}{b} + \frac{c}{d} = \frac{a}{b} \cdot 1 + 1 \cdot \frac{c}{d} = \left(\frac{a}{b}\right) \cdot \left(\frac{d}{d}\right) + \left(\frac{b}{b}\right) \cdot \left(\frac{c}{d}\right) = \frac{ad}{bd} + \frac{bc}{bd} = \frac{ad + bc}{bd}$$

Case 2: Different Denominators:

For fractions with different denominators, say $\frac{a}{b}$ and $\frac{c}{d}$, we must first alter the fractions to have a **common denominator**, say $h = bd$. We do this by multiplications by 1, where we write 1 as an appropriate fraction:

$$\frac{a}{b} + \frac{c}{d} = \frac{a}{b} \cdot 1 + 1 \cdot \frac{c}{d} = \left(\frac{a}{b}\right) \cdot \left(\frac{d}{d}\right) + \left(\frac{b}{b}\right) \cdot \left(\frac{c}{d}\right) = \frac{ad}{bd} + \frac{bc}{bd} = \frac{ad + bc}{bd}$$

We have thus derived the basic addition formula:

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd}$$

The formula is easily remembered as “cross-multiplication”:

$$\frac{a}{b} \bowtie \frac{c}{d} \rightarrow \frac{ad + bc}{bd}$$

Example 5:

$$\frac{3}{5} + \frac{7}{8} =$$

Example 5:

$$\frac{3}{5} + \frac{7}{8} = \left(\frac{3}{5}\right) \left(\frac{8}{8}\right) + \left(\frac{7}{8}\right) \left(\frac{5}{5}\right) =$$

Example 5:

$$\frac{3}{5} + \frac{7}{8} = \left(\frac{3}{5}\right) \left(\frac{8}{8}\right) + \left(\frac{7}{8}\right) \left(\frac{5}{5}\right) = \frac{24}{40} + \frac{35}{40} =$$

Example 5:

$$\frac{3}{5} + \frac{7}{8} = \left(\frac{3}{5}\right) \left(\frac{8}{8}\right) + \left(\frac{7}{8}\right) \left(\frac{5}{5}\right) = \frac{24}{40} + \frac{35}{40} = \frac{59}{40}$$

Example 6:

Example 5:

$$\frac{3}{5} + \frac{7}{8} = \left(\frac{3}{5}\right) \left(\frac{8}{8}\right) + \left(\frac{7}{8}\right) \left(\frac{5}{5}\right) = \frac{24}{40} + \frac{35}{40} = \frac{59}{40}$$

Example 6:

$$\frac{2}{3} + \frac{6}{5} =$$

Example 5:

$$\frac{3}{5} + \frac{7}{8} = \left(\frac{3}{5}\right) \left(\frac{8}{8}\right) + \left(\frac{7}{8}\right) \left(\frac{5}{5}\right) = \frac{24}{40} + \frac{35}{40} = \frac{59}{40}$$

Example 6:

$$\frac{2}{3} + \frac{6}{5} = \left(\frac{2}{3}\right) \left(\frac{5}{5}\right) + \left(\frac{6}{5}\right) \left(\frac{3}{3}\right) =$$

Example 5:

$$\frac{3}{5} + \frac{7}{8} = \left(\frac{3}{5}\right) \left(\frac{8}{8}\right) + \left(\frac{7}{8}\right) \left(\frac{5}{5}\right) = \frac{24}{40} + \frac{35}{40} = \frac{59}{40}$$

Example 6:

$$\frac{2}{3} + \frac{6}{5} = \left(\frac{2}{3}\right) \left(\frac{5}{5}\right) + \left(\frac{6}{5}\right) \left(\frac{3}{3}\right) = \frac{10}{15} + \frac{18}{15} =$$

Example 5:

$$\frac{3}{5} + \frac{7}{8} = \left(\frac{3}{5}\right) \left(\frac{8}{8}\right) + \left(\frac{7}{8}\right) \left(\frac{5}{5}\right) = \frac{24}{40} + \frac{35}{40} = \frac{59}{40}$$

Example 6:

$$\frac{2}{3} + \frac{6}{5} = \left(\frac{2}{3}\right) \left(\frac{5}{5}\right) + \left(\frac{6}{5}\right) \left(\frac{3}{3}\right) = \frac{10}{15} + \frac{18}{15} = \frac{28}{15}$$

Factor if You Can

Factor if You Can

Calculations with fractions can be simplified by factoring if there are common factors:

Example 7:

$$\frac{15}{28} + \frac{24}{35} =$$

Factor if You Can

Calculations with fractions can be simplified by factoring if there are common factors:

Example 7:

$$\frac{15}{28} + \frac{24}{35} = \frac{3 \cdot 5}{4 \cdot 7} + \frac{3 \cdot 8}{5 \cdot 7} =$$

Factor if You Can

Calculations with fractions can be simplified by factoring if there are common factors:

Example 7:

$$\frac{15}{28} + \frac{24}{35} = \frac{3 \cdot 5}{4 \cdot 7} + \frac{3 \cdot 8}{5 \cdot 7} = \left(\frac{3}{7}\right) \left(\frac{5}{4}\right) + \left(\frac{3}{7}\right) \left(\frac{8}{5}\right) =$$

Factor if You Can

Calculations with fractions can be simplified by factoring if there are common factors:

Example 7:

$$\frac{15}{28} + \frac{24}{35} = \frac{3 \cdot 5}{4 \cdot 7} + \frac{3 \cdot 8}{5 \cdot 7} = \left(\frac{3}{7}\right) \left(\frac{5}{4}\right) + \left(\frac{3}{7}\right) \left(\frac{8}{5}\right) =$$
$$\left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) =$$

Factor if You Can

Calculations with fractions can be simplified by factoring if there are common factors:

Example 7:

$$\frac{15}{28} + \frac{24}{35} = \frac{3 \cdot 5}{4 \cdot 7} + \frac{3 \cdot 8}{5 \cdot 7} = \left(\frac{3}{7}\right) \left(\frac{5}{4}\right) + \left(\frac{3}{7}\right) \left(\frac{8}{5}\right) =$$

$$\left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{5 \cdot 5 + 8 \cdot 4}{4 \cdot 5}\right) =$$

Factor if You Can

Calculations with fractions can be simplified by factoring if there are common factors:

Example 7:

$$\frac{15}{28} + \frac{24}{35} = \frac{3 \cdot 5}{4 \cdot 7} + \frac{3 \cdot 8}{5 \cdot 7} = \left(\frac{3}{7}\right) \left(\frac{5}{4}\right) + \left(\frac{3}{7}\right) \left(\frac{8}{5}\right) =$$

$$\left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{5 \cdot 5 + 8 \cdot 4}{4 \cdot 5}\right) =$$

$$\left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) =$$

Factor if You Can

Calculations with fractions can be simplified by factoring if there are common factors:

Example 7:

$$\frac{15}{28} + \frac{24}{35} = \frac{3 \cdot 5}{4 \cdot 7} + \frac{3 \cdot 8}{5 \cdot 7} = \left(\frac{3}{7}\right) \left(\frac{5}{4}\right) + \left(\frac{3}{7}\right) \left(\frac{8}{5}\right) =$$

$$\left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{5 \cdot 5 + 8 \cdot 4}{4 \cdot 5}\right) =$$

$$\left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) =$$

Factor if You Can

Calculations with fractions can be simplified by factoring if there are common factors:

Example 7:

$$\frac{15}{28} + \frac{24}{35} = \frac{3 \cdot 5}{4 \cdot 7} + \frac{3 \cdot 8}{5 \cdot 7} = \left(\frac{3}{7}\right) \left(\frac{5}{4}\right) + \left(\frac{3}{7}\right) \left(\frac{8}{5}\right) =$$

$$\left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{5 \cdot 5 + 8 \cdot 4}{4 \cdot 5}\right) =$$

$$\left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) = \frac{171}{140}$$

Factor if You Can

Calculations with fractions can be simplified by factoring if there are common factors:

Example 7:

$$\frac{15}{28} + \frac{24}{35} = \frac{3 \cdot 5}{4 \cdot 7} + \frac{3 \cdot 8}{5 \cdot 7} = \left(\frac{3}{7}\right) \left(\frac{5}{4}\right) + \left(\frac{3}{7}\right) \left(\frac{8}{5}\right) =$$

$$\left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{5 \cdot 5 + 8 \cdot 4}{4 \cdot 5}\right) =$$

$$\left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) = \frac{171}{140}$$

which would normally be done with fewer steps displayed:

Arithmetic of Fractions-10

$$\frac{15}{28} + \frac{24}{35} =$$

Arithmetic of Fractions-10

$$\frac{15}{28} + \frac{24}{35} = \left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) =$$

Arithmetic of Fractions-10

$$\frac{15}{28} + \frac{24}{35} = \left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) =$$

Arithmetic of Fractions-10

$$\frac{15}{28} + \frac{24}{35} = \left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) =$$

Arithmetic of Fractions-10

$$\frac{15}{28} + \frac{24}{35} = \left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) =$$
$$\frac{171}{140}$$

Arithmetic of Fractions-10

$$\frac{15}{28} + \frac{24}{35} = \left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) =$$
$$\frac{171}{140}$$

Note that the straightforward approach is much more cumbersome:

$$\frac{15}{28} + \frac{24}{35} =$$

$$\frac{15}{28} + \frac{24}{35} = \left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) =$$

$$\frac{171}{140}$$

Note that the straightforward approach is much more cumbersome:

$$\frac{15}{28} + \frac{24}{35} = \frac{(15)(35) + (28)(24)}{(28)(35)} =$$

$$\frac{15}{28} + \frac{24}{35} = \left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) =$$

$$\frac{171}{140}$$

Note that the straightforward approach is much more cumbersome:

$$\frac{15}{28} + \frac{24}{35} = \frac{(15)(35) + (28)(24)}{(28)(35)} = \frac{525 + 672}{980} =$$

Arithmetic of Fractions-10

$$\frac{15}{28} + \frac{24}{35} = \left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) =$$

$\frac{171}{140}$

Note that the straightforward approach is much more cumbersome:

$$\frac{15}{28} + \frac{24}{35} = \frac{(15)(35) + (28)(24)}{(28)(35)} = \frac{525 + 672}{980} = \frac{525 + 672}{980} =$$

$$\frac{15}{28} + \frac{24}{35} = \left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) =$$

$$\frac{171}{140}$$

Note that the straightforward approach is much more cumbersome:

$$\frac{15}{28} + \frac{24}{35} = \frac{(15)(35) + (28)(24)}{(28)(35)} = \frac{525 + 672}{980} = \frac{525 + 672}{980} =$$

$$\frac{1197}{980} =$$

$$\frac{15}{28} + \frac{24}{35} = \left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) =$$

$$\frac{171}{140}$$

Note that the straightforward approach is much more cumbersome:

$$\frac{15}{28} + \frac{24}{35} = \frac{(15)(35) + (28)(24)}{(28)(35)} = \frac{525 + 672}{980} = \frac{525 + 672}{980} =$$

$$\frac{1197}{980} = \frac{7(171)}{7(140)} =$$

$$\frac{15}{28} + \frac{24}{35} = \left(\frac{3}{7}\right) \left(\frac{5}{4} + \frac{8}{5}\right) = \left(\frac{3}{7}\right) \left(\frac{25 + 32}{20}\right) = \left(\frac{3}{7}\right) \left(\frac{57}{20}\right) =$$

$$\frac{171}{140}$$

Note that the straightforward approach is much more cumbersome:

$$\frac{15}{28} + \frac{24}{35} = \frac{(15)(35) + (28)(24)}{(28)(35)} = \frac{525 + 672}{980} = \frac{525 + 672}{980} =$$

$$\frac{1197}{980} = \frac{7(171)}{7(140)} = \frac{171}{140}$$

and this only works if we know that 7 is a common factor of 1197 and 980. Not many of us have this information at our finger tips!

Danger: Notice that $\frac{a}{b} + \frac{c}{d}$ does **NOT** equal

Danger: Notice that $\frac{a}{b} + \frac{c}{d}$ does **NOT** equal $\frac{a+c}{b+d}$!!

Danger: Notice that $\frac{a}{b} + \frac{c}{d}$ does **NOT** equal $\frac{a+c}{b+d}$!!

If in doubt, try $\frac{1}{2} + \frac{1}{2} \stackrel{?}{=} \frac{1+1}{2+2} = \frac{2}{4} = \frac{1}{2}$, and you know that two halves make a whole, not a half!

Danger: Notice that $\frac{a}{b} + \frac{c}{d}$ does **NOT** equal $\frac{a+c}{b+d}$!!

If in doubt, try $\frac{1}{2} + \frac{1}{2} \stackrel{?}{=} \frac{1+1}{2+2} = \frac{2}{4} = \frac{1}{2}$, and you know that two halves make a whole, not a half!

As a special case, consider a fraction $\frac{c}{d}$ added to an arbitrary number x , which we write as the fraction $\frac{x}{1}$. Then

$$x + \frac{c}{d} =$$

As a special case, consider a fraction $\frac{c}{d}$ added to an arbitrary number x , which we write as the fraction $\frac{x}{1}$. Then

$$x + \frac{c}{d} = \frac{x}{1} + \frac{c}{d} =$$

As a special case, consider a fraction $\frac{c}{d}$ added to an arbitrary number x , which we write as the fraction $\frac{x}{1}$. Then

$$x + \frac{c}{d} = \frac{x}{1} + \frac{c}{d} = \frac{xd + 1c}{1d} =$$

As a special case, consider a fraction $\frac{c}{d}$ added to an arbitrary number x , which we write as the fraction $\frac{x}{1}$. Then

$$x + \frac{c}{d} = \frac{x}{1} + \frac{c}{d} = \frac{xd + 1c}{1d} = \frac{xd + c}{d}$$

Example 8: Combine $\frac{5}{4} + 3$ into one fraction.

Example 8: Combine $\frac{5}{4} + 3$ into one fraction.

Solution

$$\frac{5}{4} + 3 =$$

Example 8: Combine $\frac{5}{4} + 3$ into one fraction.

Solution

$$\frac{5}{4} + 3 = \frac{5}{4} + 3\frac{4}{4} =$$

Example 8: Combine $\frac{5}{4} + 3$ into one fraction.

Solution

$$\frac{5}{4} + 3 = \frac{5}{4} + 3\frac{4}{4} = \frac{5}{4} + \frac{12}{4} =$$

Example 8: Combine $\frac{5}{4} + 3$ into one fraction.

Solution

$$\frac{5}{4} + 3 = \frac{5}{4} + 3\frac{4}{4} = \frac{5}{4} + \frac{12}{4} =$$

$$\frac{17}{4}$$

Subtraction

If you know how to add fractions then you know how to subtract them; the trick is to convert $-\left(\frac{c}{d}\right)$ into $\frac{-c}{d}$:

$$\frac{a}{b} - \frac{c}{d} =$$

Subtraction

If you know how to add fractions then you know how to subtract them; the trick is to convert $-\left(\frac{c}{d}\right)$ into $\frac{-c}{d}$:

$$\frac{a}{b} - \frac{c}{d} = \frac{a}{b} + \frac{-c}{d} =$$

Subtraction

If you know how to add fractions then you know how to subtract them; the trick is to convert $-\left(\frac{c}{d}\right)$ into $\frac{-c}{d}$:

$$\frac{a}{b} - \frac{c}{d} = \frac{a}{b} + \frac{-c}{d} = \frac{ad + b(-c)}{bd} =$$

Subtraction

If you know how to add fractions then you know how to subtract them; the trick is to convert $-\left(\frac{c}{d}\right)$ into $\frac{-c}{d}$:

$$\frac{a}{b} - \frac{c}{d} = \frac{a}{b} + \frac{-c}{d} = \frac{ad + b(-c)}{bd} = \frac{ad - bc}{bd},$$

so we have the subtraction formula

Subtraction

If you know how to add fractions then you know how to subtract them; the trick is to convert $-\left(\frac{c}{d}\right)$ into $\frac{-c}{d}$:

$$\frac{a}{b} - \frac{c}{d} = \frac{a}{b} + \frac{-c}{d} = \frac{ad + b(-c)}{bd} = \frac{ad - bc}{bd},$$

so we have the subtraction formula

$$\frac{a}{b} - \frac{c}{d} = \frac{ad - bc}{bd}$$

Example 9:

$$\frac{5}{7} - \frac{3}{4} =$$

Subtraction

If you know how to add fractions then you know how to subtract them; the trick is to convert $-\left(\frac{c}{d}\right)$ into $\frac{-c}{d}$:

$$\frac{a}{b} - \frac{c}{d} = \frac{a}{b} + \frac{-c}{d} = \frac{ad + b(-c)}{bd} = \frac{ad - bc}{bd},$$

so we have the subtraction formula

$$\frac{a}{b} - \frac{c}{d} = \frac{ad - bc}{bd}$$

Example 9:

$$\frac{5}{7} - \frac{3}{4} = \frac{5}{7} + \frac{-3}{4} =$$

Subtraction

If you know how to add fractions then you know how to subtract them; the trick is to convert $-\left(\frac{c}{d}\right)$ into $\frac{-c}{d}$:

$$\frac{a}{b} - \frac{c}{d} = \frac{a}{b} + \frac{-c}{d} = \frac{ad + b(-c)}{bd} = \frac{ad - bc}{bd},$$

so we have the subtraction formula

$$\frac{a}{b} - \frac{c}{d} = \frac{ad - bc}{bd}$$

Example 9:

$$\frac{5}{7} - \frac{3}{4} = \frac{5}{7} + \frac{-3}{4} = \frac{5(4) + 7(-3)}{7(4)} =$$

Subtraction

If you know how to add fractions then you know how to subtract them; the trick is to convert $-\left(\frac{c}{d}\right)$ into $\frac{-c}{d}$:

$$\frac{a}{b} - \frac{c}{d} = \frac{a}{b} + \frac{-c}{d} = \frac{ad + b(-c)}{bd} = \frac{ad - bc}{bd},$$

so we have the subtraction formula

$$\frac{a}{b} - \frac{c}{d} = \frac{ad - bc}{bd}$$

Example 9:

$$\frac{5}{7} - \frac{3}{4} = \frac{5}{7} + \frac{-3}{4} = \frac{5(4) + 7(-3)}{7(4)} = \frac{20 - 21}{28} =$$

Subtraction

If you know how to add fractions then you know how to subtract them; the trick is to convert $-\left(\frac{c}{d}\right)$ into $\frac{-c}{d}$:

$$\frac{a}{b} - \frac{c}{d} = \frac{a}{b} + \frac{-c}{d} = \frac{ad + b(-c)}{bd} = \frac{ad - bc}{bd},$$

so we have the subtraction formula

$$\frac{a}{b} - \frac{c}{d} = \frac{ad - bc}{bd}$$

Example 9:

$$\frac{5}{7} - \frac{3}{4} = \frac{5}{7} + \frac{-3}{4} = \frac{5(4) + 7(-3)}{7(4)} = \frac{20 - 21}{28} = -\frac{1}{28}$$

Factorization can again be used to simplify subtraction calculations:

Example 10:

$$\frac{5}{12} - \frac{3}{28} =$$

Factorization can again be used to simplify subtraction calculations:

Example 10:

$$\frac{5}{12} - \frac{3}{28} = \frac{5}{3(4)} - \frac{3}{4(7)} =$$

Factorization can again be used to simplify subtraction calculations:

Example 10:

$$\frac{5}{12} - \frac{3}{28} = \frac{5}{3(4)} - \frac{3}{4(7)} = \frac{1}{4} \left(\frac{5}{3} - \frac{3}{7} \right) =$$

Factorization can again be used to simplify subtraction calculations:

Example 10:

$$\frac{5}{12} - \frac{3}{28} = \frac{5}{3(4)} - \frac{3}{4(7)} = \frac{1}{4} \left(\frac{5}{3} - \frac{3}{7} \right) = \frac{1}{4} \left(\frac{5(7) - 3(3)}{3(7)} \right) =$$

Factorization can again be used to simplify subtraction calculations:

Example 10:

$$\frac{5}{12} - \frac{3}{28} = \frac{5}{3(4)} - \frac{3}{4(7)} = \frac{1}{4} \left(\frac{5}{3} - \frac{3}{7} \right) = \frac{1}{4} \left(\frac{5(7) - 3(3)}{3(7)} \right) =$$

$$\frac{1}{4} \left(\frac{35 - 9}{21} \right) =$$

Factorization can again be used to simplify subtraction calculations:

Example 10:

$$\frac{5}{12} - \frac{3}{28} = \frac{5}{3(4)} - \frac{3}{4(7)} = \frac{1}{4} \left(\frac{5}{3} - \frac{3}{7} \right) = \frac{1}{4} \left(\frac{5(7) - 3(3)}{3(7)} \right) =$$

$$\frac{1}{4} \left(\frac{35 - 9}{21} \right) = \frac{1}{4} \left(\frac{26}{21} \right) =$$

Factorization can again be used to simplify subtraction calculations:

Example 10:

$$\frac{5}{12} - \frac{3}{28} = \frac{5}{3(4)} - \frac{3}{4(7)} = \frac{1}{4} \left(\frac{5}{3} - \frac{3}{7} \right) = \frac{1}{4} \left(\frac{5(7) - 3(3)}{3(7)} \right) =$$

$$\frac{1}{4} \left(\frac{35 - 9}{21} \right) = \frac{1}{4} \left(\frac{26}{21} \right) = \frac{1}{2} \left(\frac{13}{21} \right) =$$

Factorization can again be used to simplify subtraction calculations:

Example 10:

$$\frac{5}{12} - \frac{3}{28} = \frac{5}{3(4)} - \frac{3}{4(7)} = \frac{1}{4} \left(\frac{5}{3} - \frac{3}{7} \right) = \frac{1}{4} \left(\frac{5(7) - 3(3)}{3(7)} \right) =$$

$$\frac{1}{4} \left(\frac{35 - 9}{21} \right) = \frac{1}{4} \left(\frac{26}{21} \right) = \frac{1}{2} \left(\frac{13}{21} \right) = \frac{13}{42}$$

Division

To divide $\frac{a}{b}$ by $\frac{c}{d}$, we remember that $\frac{c}{d} \cdot \frac{d}{c} = \frac{cd}{cd} = 1$,

and multiply by 1 in a special form:

$$\frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} =$$

Division

To divide $\frac{a}{b}$ by $\frac{c}{d}$, we remember that $\frac{c}{d} \cdot \frac{d}{c} = \frac{cd}{cd} = 1$,

and multiply by 1 in a special form:

$$\frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} \cdot \frac{\left(\frac{d}{c}\right)}{\left(\frac{d}{c}\right)} =$$

Division

To divide $\frac{a}{b}$ by $\frac{c}{d}$, we remember that $\frac{c}{d} \cdot \frac{d}{c} = \frac{cd}{cd} = 1$,

and multiply by 1 in a special form:

$$\frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} \cdot \frac{\left(\frac{d}{c}\right)}{\left(\frac{d}{c}\right)} = \frac{\left(\frac{a}{b}\right)\left(\frac{d}{c}\right)}{\left(\frac{c}{d}\right)\left(\frac{d}{c}\right)} =$$

Division

To divide $\frac{a}{b}$ by $\frac{c}{d}$, we remember that $\frac{c}{d} \cdot \frac{d}{c} = \frac{cd}{cd} = 1$,

and multiply by 1 in a special form:

$$\frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} \cdot \frac{\left(\frac{d}{c}\right)}{\left(\frac{d}{c}\right)} = \frac{\left(\frac{a}{b}\right)\left(\frac{d}{c}\right)}{\left(\frac{c}{d}\right)\left(\frac{d}{c}\right)} = \frac{\left(\frac{ad}{bc}\right)}{1} =$$

Division

To divide $\frac{a}{b}$ by $\frac{c}{d}$, we remember that $\frac{c}{d} \cdot \frac{d}{c} = \frac{cd}{cd} = 1$,

and multiply by 1 in a special form:

$$\frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} \cdot \frac{\left(\frac{d}{c}\right)}{\left(\frac{d}{c}\right)} = \frac{\left(\frac{a}{b}\right)\left(\frac{d}{c}\right)}{\left(\frac{c}{d}\right)\left(\frac{d}{c}\right)} = \frac{\left(\frac{ad}{bc}\right)}{1} = \frac{ad}{bc}$$

Our basic division formula is thus

Division

To divide $\frac{a}{b}$ by $\frac{c}{d}$, we remember that $\frac{c}{d} \cdot \frac{d}{c} = \frac{cd}{cd} = 1$,

and multiply by 1 in a special form:

$$\frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} \cdot \frac{\left(\frac{d}{c}\right)}{\left(\frac{d}{c}\right)} = \frac{\left(\frac{a}{b}\right)\left(\frac{d}{c}\right)}{\left(\frac{c}{d}\right)\left(\frac{d}{c}\right)} = \frac{\left(\frac{ad}{bc}\right)}{1} = \frac{ad}{bc}$$

Our basic division formula is thus

$$\frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{d}\right)} = \frac{ad}{bc}$$

Example 11: $\frac{\frac{2}{3}}{\frac{5}{4}} =$

Example 11: $\frac{2}{3} \frac{3}{5} \frac{4}{4} = \left(\frac{2}{3}\right) \left(\frac{4}{5}\right) =$

Example 11: $\frac{\frac{2}{3}}{\frac{5}{4}} = \left(\frac{2}{3}\right) \left(\frac{4}{5}\right) = \frac{8}{15}$

Example 12:

$$\frac{\frac{2}{3} + \frac{4}{7}}{\frac{5}{7} + \frac{3}{4}} =$$

Example 11:

$$\frac{\frac{2}{3}}{\frac{5}{4}} = \left(\frac{2}{3}\right) \left(\frac{4}{5}\right) = \frac{8}{15}$$

Example 12:

$$\frac{\frac{2}{3} + \frac{4}{7}}{\frac{5}{7} + \frac{3}{4}} = \frac{\frac{14+12}{21}}{\frac{20+21}{28}} =$$

Example 11: $\frac{\frac{2}{3}}{\frac{5}{4}} = \left(\frac{2}{3}\right) \left(\frac{4}{5}\right) = \frac{8}{15}$

Example 12:

$$\frac{\frac{2}{3} + \frac{4}{7}}{\frac{5}{7} + \frac{3}{4}} = \frac{\frac{14+12}{21}}{\frac{20+21}{28}} = \frac{\frac{26}{21}}{\frac{41}{28}} = \left(\frac{26}{21}\right) \left(\frac{28}{41}\right) = \left(\frac{26}{3}\right) \left(\frac{4}{41}\right) =$$

Example 11: $\frac{\frac{2}{3}}{\frac{5}{4}} = \left(\frac{2}{3}\right) \left(\frac{4}{5}\right) = \frac{8}{15}$

Example 12:

$$\frac{\frac{2}{3} + \frac{4}{7}}{\frac{5}{7} + \frac{3}{4}} = \frac{\frac{14+12}{21}}{\frac{20+21}{28}} = \frac{\frac{26}{21}}{\frac{41}{28}} = \left(\frac{26}{21}\right) \left(\frac{28}{41}\right) = \left(\frac{26}{3}\right) \left(\frac{4}{41}\right) = \frac{104}{123}$$

Thus, to simplify the quotient of two fractions, we multiply numerator by the inverse of the denominator," i.e.,

$$\frac{a}{b} \div \frac{c}{d} =$$

Thus, to simplify the quotient of two fractions, we multiply numerator by the inverse of the denominator," i.e.,

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c} =$$

Thus, to simplify the quotient of two fractions, we multiply numerator by the inverse of the denominator," i.e.,

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \cdot \frac{d}{c} = \frac{ad}{bc}$$

Be careful not to confuse what is divided by what: the

expression $\frac{\frac{a}{b}}{c}$ is ambiguous:

Does it mean: $\frac{\left(\frac{a}{b}\right)}{c}$ or $\frac{a}{\left(\frac{b}{c}\right)}$?

These two expressions mean very different things:

Does it mean: $\frac{\left(\frac{a}{b}\right)}{c}$ or $\frac{a}{\left(\frac{b}{c}\right)}$?

These two expressions mean very different things:

$$\frac{\left(\frac{a}{b}\right)}{c} =$$

Does it mean: $\frac{\left(\frac{a}{b}\right)}{c}$ or $\frac{a}{\left(\frac{b}{c}\right)}$?

These two expressions mean very different things:

$$\frac{\left(\frac{a}{b}\right)}{c} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{1}\right)} =$$

Does it mean: $\frac{\left(\frac{a}{b}\right)}{c}$ or $\frac{a}{\left(\frac{b}{c}\right)}$?

These two expressions mean very different things:

$$\frac{\left(\frac{a}{b}\right)}{c} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{1}\right)} = \left(\frac{a}{b}\right) \left(\frac{1}{c}\right) =$$

Does it mean: $\frac{\left(\frac{a}{b}\right)}{c}$ or $\frac{a}{\left(\frac{b}{c}\right)}$?

These two expressions mean very different things:

$$\frac{\left(\frac{a}{b}\right)}{c} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{1}\right)} = \left(\frac{a}{b}\right) \left(\frac{1}{c}\right) = \frac{a \cdot 1}{bc} =$$

Does it mean: $\frac{\left(\frac{a}{b}\right)}{c}$ or $\frac{a}{\left(\frac{b}{c}\right)}$?

These two expressions mean very different things:

$$\frac{\left(\frac{a}{b}\right)}{c} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{1}\right)} = \left(\frac{a}{b}\right) \left(\frac{1}{c}\right) = \frac{a \cdot 1}{bc} = \frac{a}{bc}$$

Does it mean: $\frac{\left(\frac{a}{b}\right)}{c}$ or $\frac{a}{\left(\frac{b}{c}\right)}$?

These two expressions mean very different things:

$$\frac{\left(\frac{a}{b}\right)}{c} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{1}\right)} = \left(\frac{a}{b}\right) \left(\frac{1}{c}\right) = \frac{a \cdot 1}{bc} = \frac{a}{bc}$$

and

$$\frac{a}{\left(\frac{b}{c}\right)} =$$

Does it mean: $\frac{\left(\frac{a}{b}\right)}{c}$ or $\frac{a}{\left(\frac{b}{c}\right)}$?

These two expressions mean very different things:

$$\frac{\left(\frac{a}{b}\right)}{c} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{1}\right)} = \left(\frac{a}{b}\right) \left(\frac{1}{c}\right) = \frac{a \cdot 1}{bc} = \frac{a}{bc}$$

and

$$\frac{a}{\left(\frac{b}{c}\right)} = \frac{\left(\frac{a}{1}\right)}{\left(\frac{b}{c}\right)} =$$

Does it mean: $\frac{\left(\frac{a}{b}\right)}{c}$ or $\frac{a}{\left(\frac{b}{c}\right)}$?

These two expressions mean very different things:

$$\frac{\left(\frac{a}{b}\right)}{c} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{1}\right)} = \left(\frac{a}{b}\right) \left(\frac{1}{c}\right) = \frac{a \cdot 1}{bc} = \frac{a}{bc}$$

and

$$\frac{a}{\left(\frac{b}{c}\right)} = \frac{\left(\frac{a}{1}\right)}{\left(\frac{b}{c}\right)} = \left(\frac{a}{1}\right) \left(\frac{c}{b}\right) =$$

Does it mean: $\frac{\left(\frac{a}{b}\right)}{c}$ or $\frac{a}{\left(\frac{b}{c}\right)}$?

These two expressions mean very different things:

$$\frac{\left(\frac{a}{b}\right)}{c} = \frac{\left(\frac{a}{b}\right)}{\left(\frac{c}{1}\right)} = \left(\frac{a}{b}\right) \left(\frac{1}{c}\right) = \frac{a \cdot 1}{bc} = \frac{a}{bc}$$

and

$$\frac{a}{\left(\frac{b}{c}\right)} = \frac{\left(\frac{a}{1}\right)}{\left(\frac{b}{c}\right)} = \left(\frac{a}{1}\right) \left(\frac{c}{b}\right) = \frac{ac}{b}$$

Example 13:

$$\frac{\left(\frac{1}{2}\right)}{3} =$$

Example 13:

$$\frac{\left(\frac{1}{2}\right)}{3} = \left(\frac{1}{2}\right) \left(\frac{1}{3}\right) =$$

Example 13:

$$\frac{\binom{1}{2}}{3} = \binom{1}{2} \binom{1}{3} = \frac{1}{6} \text{ (Remember that pizza?)}$$

Example 14:

$$\frac{1}{\binom{2}{3}} =$$

Example 13:

$$\frac{\binom{1}{2}}{3} = \binom{1}{2} \binom{1}{3} = \frac{1}{6} \text{ (Remember that pizza?)}$$

Example 14:

$$\frac{1}{\binom{2}{3}} = \frac{3}{2}$$