

## Completing the Square

With quadratic polynomials it is often important to **complete the square**. As a matter of fact, the Quadratic Formula for the roots of  $ax^2 + bx + c = 0$  is derived by using this method.

The basic idea is that we rewrite  $ax^2 + bx + c$  in the form

$$a[(x + h)^2 + k].$$

**Example 1:**  $x^2 + 2x + 2 = \overbrace{x^2 + 2x + 1}^{(x+1)^2} + 1 = (x + 1)^2 + 1$

What we try to do is rearrange the expression so that its first three terms are a perfect square. An organized approach to this is to add an expression that equals 0 in just the right way:

$$x^2 + 2x + 2 = x^2 + 2x + \underbrace{(1 - 1)}_0 + 2 = \overbrace{x^2 + 2x + 1}^{(x+1)^2} + (-1) + 2 = (x + 1)^2 + 1$$

But how did we know to add 0 in the form  $1 - 1$ ? The answer is that 1 is one-half of the coefficient 2 of  $x$  in the expression  $x^2 + 2x + 2$ .

Things get more complicated when the coefficient of  $x$  isn't equal to 2:

**Example 2:**  $x^2 + 4x + 2 = x^2 + 2(2)x + \underbrace{(2^2 - 2^2)}_0 + 2 = \overbrace{x^2 + 2(2)x + 2^2}^{(x+2)^2} - 2^2 + 2 = (x + 2)^2 - 2$

Here the coefficient of  $x$  was 4, so we divided it by 2 to get 2, and then added 0 in the form  $2^2 - 2^2$ .

When we are dealing with an expression like  $x^2 + bx + c$ , we have to add 0 in the form of the square of half of  $b$  minus itself:

$$x^2 + bx + c = x^2 + 2\left(\frac{b}{2}\right)x + c = x^2 + 2\left(\frac{b}{2}\right)x + \underbrace{\left(\left(\frac{b}{2}\right)^2 - \left(\frac{b}{2}\right)^2\right)}_0 + c =$$

$$\overbrace{x^2 + 2\left(\frac{b}{2}\right)x + \left(\frac{b}{2}\right)^2}^{(x+\frac{b}{2})^2} - \left(\frac{b}{2}\right)^2 + c = \left(x + \frac{b}{2}\right)^2 - \left(\frac{b}{2}\right)^2 + c$$

### Example 3:

$$x^2 + 5x + 3 = x^2 + 2\left(\frac{5}{2}\right)x + \underbrace{\left(\left(\frac{5}{2}\right)^2 - \left(\frac{5}{2}\right)^2\right)}_0 + 3$$

$$\overbrace{x^2 + 2\left(\frac{5}{2}\right)x + \left(\frac{5}{2}\right)^2}^{(x+\frac{5}{2})^2} - \left(\frac{5}{2}\right)^2 + 3\left(\frac{4}{4}\right) = \left(x + \frac{5}{2}\right)^2 - \frac{25}{4} + \frac{12}{4} =$$

$$\left(x + \frac{5}{2}\right)^2 - \frac{13}{4}$$

### Example 4:

$$x^2 + 9x + 5 = x^2 + 2 \left(\frac{9}{2}\right) x + \underbrace{\left(\left(\frac{9}{2}\right)^2 - \left(\frac{9}{2}\right)^2\right)}_0 + 5$$

$$\overbrace{x^2 + 2 \left(\frac{9}{2}\right) x + \left(\frac{9}{2}\right)^2}^{(x+\frac{9}{2})^2} - \left(\frac{9}{2}\right)^2 + 5 \left(\frac{4}{4}\right) = \left(x + \frac{9}{2}\right)^2 - \frac{81}{4} + \frac{20}{4} =$$

$$\left(x + \frac{9}{2}\right)^2 - \frac{61}{4}$$

Things get even more complicated when the coefficient  $a$  of  $x^2$  is not equal to 1: we have to factor it out before proceeding as above:

$$ax^2 + bx + c = a \left[ x^2 + \frac{b}{a}x + \frac{c}{a} \right] =$$

$$a \left[ x^2 + 2 \left(\frac{b}{2a}\right) x + \underbrace{\left(\left(\frac{b}{2a}\right)^2 - \left(\frac{b}{2a}\right)^2\right)}_0 + \frac{c}{a} \right] =$$

$$a \left[ \overbrace{x^2 + 2 \left(\frac{b}{2a}\right) x + \left(\frac{b}{2a}\right)^2}^{(x+\frac{b}{2a})^2} - \left(\frac{b}{2a}\right)^2 + \frac{c}{a} \right] = a \left[ \left(x + \frac{b}{2a}\right)^2 - \frac{b^2}{4a^2} + \frac{4ac}{4a^2} \right] =$$

$$a \left[ \left(x + \frac{b}{2a}\right)^2 - \frac{b^2 - 4ac}{4a^2} \right]$$

### Example 5:

$$\begin{aligned}
3x^2 + 5x + 2 &= 3 \left[ x^2 + \frac{5}{3}x + \frac{2}{3} \right] = \\
3 \left[ x^2 + 2 \left( \frac{5}{6} \right) x + \underbrace{\left( \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right)}_0 + \frac{2}{3} \right] &= \\
3 \left[ x^2 + 2 \left( \frac{5}{6} \right) x + \overbrace{\left( \frac{5}{6} \right)^2}^{(x + \frac{5}{6})^2} - \left( \frac{5}{6} \right)^2 + \frac{2}{3} \left( \frac{12}{12} \right) \right] &= \\
3 \left[ \left( x + \frac{5}{6} \right)^2 - \frac{25}{36} + \frac{24}{36} \right] &= 3 \left[ \left( x + \frac{5}{6} \right)^2 - \frac{1}{36} \right] = 3 \left[ \left( x + \frac{5}{6} \right)^2 - \left( \frac{1}{6} \right)^2 \right] = \\
3 \left( x + \frac{5}{6} + \frac{1}{6} \right) \left( x + \frac{5}{6} - \frac{1}{6} \right) &= 3(x + 1) \left( x + \frac{2}{3} \right) = \boxed{(x + 1)(3x + 2)}
\end{aligned}$$


---

We return to the general situation. By completing squares, we derived the equation:

$$ax^2 + bx + c = a \left[ \left( x + \frac{b}{2a} \right)^2 - \frac{b^2 - 4ac}{4a^2} \right]$$

We use this to solve the equation  $ax^2 + bx + c = 0$ : since  $a \neq 0$  we have

$$\begin{aligned}
ax^2 + bx + c &= 0 && \Leftrightarrow \\
\left( x + \frac{b}{2a} \right)^2 - \frac{b^2 - 4ac}{4a^2} &= 0 && \Leftrightarrow \\
\left( x + \frac{b}{2a} \right)^2 &= \frac{b^2 - 4ac}{4a^2} && \Leftrightarrow (\text{if } b^2 - 4ac \geq 0) \\
x + \frac{b}{2a} &= \pm \sqrt{\frac{b^2 - 4ac}{4a^2}} = \pm \frac{\sqrt{b^2 - 4ac}}{2a} && \Leftrightarrow \\
x &= -\frac{b}{2a} \pm \frac{\sqrt{b^2 - 4ac}}{2a} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}, \text{ which is the familiar Quadratic Formula for the roots of the general quadratic equation.}
\end{aligned}$$