

## Limit Laws

We let  $\lim$  stand for any one of the limits  $\lim_{x \rightarrow a}$ ,  $\lim_{x \rightarrow a^-}$ ,  $\lim_{x \rightarrow a^+}$ ,  $\lim_{x \rightarrow -\infty}$ , or  $\lim_{x \rightarrow +\infty}$ , and suppose  $c$  is a constant and that  $\lim f(x)$  and  $\lim g(x)$  exist. Then

$$(1) \lim [f(x) + g(x)] = \lim f(x) + \lim g(x)$$

$$(2) \lim [f(x) - g(x)] = \lim f(x) - \lim g(x)$$

$$(3) \lim [f(x)g(x)] = \lim f(x) \lim g(x)$$

$$(4) \lim \frac{f(x)}{g(x)} = \frac{\lim f(x)}{\lim g(x)} \text{ if } \lim g(x) \text{ is not equal to } 0.$$

$$(5) \lim \sqrt[n]{f(x)} = \sqrt[n]{\lim f(x)}, \text{ if } n \text{ is a positive integer and } \lim f(x) \geq 0 \text{ is also required if } n \text{ is even.}$$

$$(6) \text{ If } n \text{ is a positive integer, } \lim (f(x))^n = (\lim f(x))^n$$

$$(7) \lim cf(x) = c \lim f(x)$$

$$(8) \text{ If } c \text{ is a constant then } \lim c = c \text{ for any } a.$$

---

The next three limit laws do not apply to limits as  $x \rightarrow \pm\infty$ :

$$(9) \lim_{x \rightarrow a} x = a, \lim_{x \rightarrow a^-} x = a, \lim_{x \rightarrow a^+} x = a, \text{ for any } a.$$

$$(10) \lim_{x \rightarrow a} x^n = a^n, \lim_{x \rightarrow a^-} x^n = a^n, \lim_{x \rightarrow a^+} x^n = a^n, \text{ for any } a, \text{ if } n \text{ is a positive integer.}$$

$$(11) \lim_{x \rightarrow a} \sqrt[n]{x} = \sqrt[n]{a}, \lim_{x \rightarrow a^-} \sqrt[n]{x} = \sqrt[n]{a}, \lim_{x \rightarrow a^+} \sqrt[n]{x} = \sqrt[n]{a}, \text{ for any } a > 0,$$

if  $n$  is a positive integer.

(12) If  $f(x) = g(x)$  for all  $x$  except possibly  $x = a$ , then  $\lim_{x \rightarrow a} f(x) = \lim_{x \rightarrow a} g(x)$  if either limit exists.

(13) If  $f(x)$  is a polynomial or a rational function and  $a$  is in the domain of  $f$ , then  $\lim_{x \rightarrow a} f(x) = f(a)$

### Examples:

**Example 1:**  $\lim_{x \rightarrow -1} [x^5 - 3x^3 + 1](x^2 - 2) = [(-1)^5 - 3(-1)^3 + 1]((-1)^2 - 2) =$

$$[-1 - 3(-1) + 1](1 - 2) = [-1 + 3 + 1](-1) = (3)(-1) = -3$$

### Example 2:

$$\lim_{x \rightarrow 25} \frac{\sqrt{x}}{x + 25} = \frac{\sqrt{25}}{25 + 25} = \frac{5}{50} = \frac{1}{10}$$

### Example 3:

$$\lim_{x \rightarrow 9} \frac{\sqrt{x} - 3}{x - 9} = \lim_{x \rightarrow 9} \left( \frac{\sqrt{x} - 3}{x - 9} \right) \left( \frac{\sqrt{x} + 3}{\sqrt{x} + 3} \right) = \lim_{x \rightarrow 9} \frac{(\sqrt{x})^2 - 3^2}{(x - 9)(\sqrt{x} + 3)} =$$

$$\lim_{x \rightarrow 9} \frac{x - 9}{(x - 9)(\sqrt{x} + 3)} = \lim_{x \rightarrow 9} \frac{1}{\sqrt{x} + 3} = \frac{1}{\lim_{x \rightarrow 9} (\sqrt{x} + 3)} = \frac{1}{\sqrt{9} + 3} = \frac{1}{3 + 3} = \frac{1}{6}$$

### Example 4:

$$\lim_{x \rightarrow 2} \frac{x - 2}{x^2 - 4} = \lim_{x \rightarrow 2} \frac{x - 2}{(x - 2)(x + 2)} = \lim_{x \rightarrow 2} \frac{1}{x + 2} = \frac{1}{2 + 2} = \frac{1}{4}$$

### Example 5:

$$\lim_{x \rightarrow 1} \frac{x^2 - 1}{x - 1} = \lim_{x \rightarrow 1} \frac{(x + 1)(x - 1)}{x - 1} = \lim_{x \rightarrow 1} (x + 1) = 1 + 1 = 2$$

---

### Example 6: $\lim_{x \rightarrow 0} |x|$

**DANGER!**  $|x|$  is a function defined by a multicolumn formula, and different formulas apply for  $|x|$  on either side of the limiting value of  $x$ , namely 0. We must calculate the two one-sided limits:

$$\lim_{x \rightarrow 0^-} |x| = \lim_{x \rightarrow 0^-} (-x) = (-0) = 0$$

$$\lim_{x \rightarrow 0^+} |x| = \lim_{x \rightarrow 0^+} x = 0$$

Since the left and right hand limits are equal, we can conclude that the two-sided limit does exist and equals the common value 0.

---

### Example 7:

$$\lim_{x \rightarrow 0} \frac{|x|}{x}$$


Note that the limiting value of  $x$ , 0, is not in the domain of the function  $\frac{|x|}{x}$ .

Again,  $|x|$  is a function defined by a multicolumn formula, and different formulas apply for  $x$  on either side of the limiting value of  $x$ , so again we must calculate the two one-sided limits:

$$\lim_{x \rightarrow 0^-} \frac{|x|}{x} = \lim_{x \rightarrow 0^-} \frac{-x}{x} = \lim_{x \rightarrow 0^-} -1 = -1$$

$$\lim_{x \rightarrow 0^+} \frac{|x|}{x} = \lim_{x \rightarrow 0^+} \frac{x}{x} = \lim_{x \rightarrow 0^+} 1 = 1$$

Since the left and right hand limits are unequal, we can conclude that the two-sided limit does not exist.



## Limits of Rational Functions

If  $f(x) = \frac{p(x)}{q(x)}$  where  $p(x)$  and  $q(x)$  are polynomials, then

$$\lim_{x \rightarrow a} f(x) = f(a) = \frac{p(a)}{q(a)} \quad \text{if } q(a) \neq 0$$

If  $q(a) = 0$ , then there are a number of possibilities:

if  $p(a) \neq 0$ , we could have  $\lim_{x \rightarrow a} f(x) = +\infty$ , or  $\lim_{x \rightarrow a} f(x) = -\infty$ , or  
 $\lim_{x \rightarrow a^-} f(x) = -\infty$  and  $\lim_{x \rightarrow a^+} f(x) = +\infty$ , or  $\lim_{x \rightarrow a^-} f(x) = +\infty$  and  $\lim_{x \rightarrow a^+} f(x) = -\infty$ .

If  $p(a) = 0$ , it is possible, but not guaranteed, that  $\lim_{x \rightarrow a} f(x)$  can exist.

**Example 8:**  $\lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2}$ .

Here  $a = 2$ ,  $p(x) = x^2 - 4$ ,  $q(x) = x - 2$ , so  $p(a) = p(2) = 0$ ,  $q(a) = q(2) = 0$ . We have to do some algebraic manipulation to compute the limit:

$$\lim_{x \rightarrow 2} \frac{x^2 - 4}{x - 2} = \lim_{x \rightarrow 2} \frac{(x - 2)(x + 2)}{x - 2} = \lim_{x \rightarrow 2} x + 2 = 2 + 2 = 4.$$

## Rational Functions at $\pm \infty$

**Theorem:** If  $f(x) = \frac{p(x)}{q(x)}$  where  $p(x)$  and  $q(x)$  are polynomials, then:

(1) if the degree of  $p(x)$  is less than the degree of  $q(x)$ ,  $\lim_{x \rightarrow \pm \infty} f(x) = 0$ ,

(2) if the degree of  $p(x)$  is greater than the degree of  $q(x)$ ,  $\lim_{x \rightarrow \pm \infty} f(x) = \pm \infty$ ,

(3) if the degree of  $p(x)$  equals the degree of  $q(x)$ ,  $\lim_{x \rightarrow \pm \infty} f(x) = \frac{a_0}{b_0}$ ,  
where  $a_0$  and  $b_0$  are the leading coefficients of  $p(x)$  and  $q(x)$ .

**Example 9:** The **Floor and Ceiling Functions** .

We define  $\lfloor x \rfloor$ , the **floor of  $x$** , to be the largest integer that is less than or equal to  $x$ , and  $\lceil x \rceil$ , the **ceiling of  $x$** , to be the least integer that is greater than or equal to  $x$ . In other words, the floor function rounds  $x$  down to the nearest integer and the ceiling function rounds  $x$  up to the nearest integer. They are both examples of step functions. They both have two-sided limits at non-integer values:

If  $a$  is not an integer, then  $\lim_{x \rightarrow a^-} \lfloor x \rfloor = \lfloor a \rfloor$ , and  $\lim_{x \rightarrow a^-} \lceil x \rceil = \lceil a \rceil$ , but if  $a$  is an integer, things are different:

$$\lim_{x \rightarrow a^-} \lfloor x \rfloor = \lfloor a \rfloor - 1,$$

$$\lim_{x \rightarrow a^+} \lfloor x \rfloor = \lfloor a \rfloor$$

and

$$\lim_{x \rightarrow a^-} \lceil x \rceil = \lceil a \rceil,$$

$$\lim_{x \rightarrow a^+} \lceil x \rceil = \lceil a \rceil + 1$$

**The Floor Function:**

**The Ceiling Function:**

