

ON THE REAL MULTIDIMENSIONAL RATIONAL K -MOMENT PROBLEM

JAKA CIMPRIC, MURRAY MARSHALL, AND TIM NETZER

ABSTRACT. We present a solution to the real multidimensional rational K -moment problem, where K is defined by finitely many polynomial inequalities. More precisely, let S be a finite set of real polynomials in $\underline{X} = (X_1, \dots, X_n)$ such that the corresponding basic closed semialgebraic set K_S is nonempty. Let $E = D^{-1}\mathbb{R}[\underline{X}]$ be a localization of the real polynomial algebra, and T_S^E the preordering on E generated by S . We show that every linear functional L on E such that $L(T_S^E) \geq 0$ is represented by a positive measure μ on a certain subset of K_S , provided D contains an element that grows fast enough on K_S .

1. INTRODUCTION

The *moment problem* for a commutative unital \mathbb{R} -algebra E asks to characterize real positive linear functionals on E which can be represented as integrals over measures on an appropriate representation space of E . If supports of the measures are required to lie in a prescribed subset K of the representation space then we talk about the *K -moment problem* on E .

A solution to the K -moment problem on $\mathbb{R}[\underline{X}] = \mathbb{R}[X_1, \dots, X_n]$ for a compact basic closed semialgebraic set K was given by K. Schmüdgen in [21]. The aim of this paper is to extend his result, both in the compact and non-compact case, to localizations of the polynomial algebra, i.e. to algebras of the form $E = D^{-1}\mathbb{R}[\underline{X}]$, where D is a multiplicative set. The case $n = 1$, D generated by $X_1 - \alpha$ for countably many real α , and K a compact basic closed semialgebraic set has already been done by J. D. Chandler in [8]. Several papers deal with the case $n = 1$, D generated by $X_1 - \alpha$ for countably many real α and $K = \mathbb{R}$, see [6, 7] for surveys. In this case the existence of the solution is rather trivial and the emphasis is on the uniqueness of solutions. The multi-dimensional case with D generated by $1 + \sum_{i=1}^n X_i^2 + \sum_{i=1}^m g_i^2$ where

Date: July 30, 2008 and, in revised form, October 9th, 2009.

2000 Mathematics Subject Classification. 44A60, 14P99.

Key words and phrases. moment problem, positive polynomials, sums of squares.

$g_1 \geq 0, \dots, g_m \geq 0$ are the defining relations of K has been done by M. Putinar and F.-H. Vasilescu in [16].

To describe our main results we need some terminology. For a finite subset $S = \{g_1, \dots, g_m\}$ of $\mathbb{R}[\underline{X}]$ write $K_S = \{a \in \mathbb{R}^n \mid g_1(a) \geq 0, \dots, g_m(a) \geq 0\}$ and T_S^E for the set of all finite sums of elements of the form $e^2 g_1^{\nu_1} \cdots g_m^{\nu_m}$ where $e \in E = D^{-1}\mathbb{R}[\underline{X}]$ and $\nu_1, \dots, \nu_m \in \{0, 1\}$. We say that a rational function $R \in E$ is $\succeq 0$ on a set $\mathcal{X} \subseteq \mathbb{R}^n$, if there exist $f \in \mathbb{R}[\underline{X}]$ and $d \in D$ such that $R = \frac{f}{d}$ and $fd \geq 0$ on \mathcal{X} . We write $\mathcal{Z}(d)$ for the set of real zeros of a polynomial d .

Our main results can be summarized as follows:

Theorem. *Let S be a finite subset of $\mathbb{R}[\underline{X}]$, D a multiplicative subset of $\mathbb{R}[\underline{X}]$ such that $1 \in D$, $0 \notin D$, and let R be an element of $E = D^{-1}\mathbb{R}[\underline{X}]$. Assume there exists an element $p \in D$ such that $p \geq 1$ on K_S and $kp \geq \sum_{i=1}^n X_i^2$ on K_S for some integer $k \geq 1$. Then*

- (1) *R belongs to the closure of T_S^E in the finest locally convex topology on E if and only if $R \succeq 0$ on $K_S \setminus \bigcup_{d \in D} \mathcal{Z}(d)$,*
- (2) *R belongs to the closure of T_S^E in the topology of finitely open sets on E if and only if $R \succeq 0$ on K_S .*
- (3) *For every linear functional L on E such that $L(T_S^E) \geq 0$, there exists a measure μ on $\overline{K_S \setminus \bigcup_{d \in D} \mathcal{Z}(d)}$ such that*

$$L\left(\frac{f}{d}\right) = \int \frac{f}{d} d\mu \quad \text{for every } \frac{f}{d} \in E.$$

See Theorems 1, 2 and 4. These results carry over to localizations of an arbitrary finitely generated \mathbb{R} -algebra; see Theorem 5. Assertion (3) of the theorem solves the moment problem on E . For the proof of assertion (3) we use assertion (1) together with a certain rational version of the Riesz-Haviland Theorem which is a generalization of [11, Theorem 3.2.2] and [24, Theorem 3.2]; see Theorem 3.

The conditions on the polynomial p (that $p \geq 1$ on K_S and $kp \geq \sum_{i=1}^n X_i^2$ on K_S for some integer $k \geq 1$) are paraphrased by saying that “ p grows fast enough on K_S ”. See also [11, Theorem 6.2.3] and [22, Theorem 5.1]. One can always take $p = 1 + \sum_{i=1}^n X_i^2$ (provided, of course, that $1 + \sum_{i=1}^n X_i^2 \in D$). If K_S is compact one can take $p = 1$, so, in this case, the only assumptions on the multiplicative set D are the trivial ones $1 \in D, 0 \notin D$.

2. PREORDERINGS AND THE FINEST LOCALLY CONVEX TOPOLOGY

Definition. Let E be an \mathbb{R} -vector space. A set $U \subseteq E$ is called *absorbent*, if for every $x \in E$ there exists $\lambda > 0$ such that $x \in \lambda U$. U is called *symmetric*, if $\lambda U \subseteq U$ for all $\lambda \in [-1, 1]$.

The set of all convex, absorbent and symmetric subsets of E forms a zero neighborhood base of a vector space topology on E (see [5, II.25]). It is called the *finest locally convex topology* on E and the collection of all open sets is denoted by \mathcal{T}_ω .

The following are well-known:

- (T1) (E, \mathcal{T}_ω) is a topological vector space, i.e. addition and scalar multiplication are continuous. Moreover, (E, \mathcal{T}_ω) is Hausdorff.
- (T2) Every linear mapping from (E, \mathcal{T}_ω) to a vector space with any locally convex topology is continuous. In particular, all functionals on (E, \mathcal{T}_ω) are continuous.

These properties are proved in [5, II.26] .

A subset T of a commutative ring E with 1 is a *preordering* if it is closed under addition and multiplication and if it contains the set $E^2 = \{a^2 \mid a \in E\}$. As usual, $\sum E^2$ denotes the preordering of E consisting of sums of squares.

Lemma 1. *For any preordering T on any commutative \mathbb{R} -algebra E , its closure in \mathcal{T}_ω is also a preordering.*

Proof. Since $T \subseteq \overline{T}$, $E^2 \subseteq \overline{T}$. Since the addition on E is continuous, it follows that \overline{T} is closed under addition. We can't use the same argument for multiplication because by [25, p. 734, Theorem 2], the multiplication on $E = \mathbb{C}(X)$ is not continuous. However, for every $f \in E$, the mapping $\phi_f: E \rightarrow E$, $\phi_f(h) = fh$, is linear, hence continuous by (T2). So, for every $f \in T$,

$$\phi_f(\overline{T}) \subseteq \overline{\phi_f(T)} \subseteq \overline{T}$$

holds, as $\phi_f(T) \subseteq T$. It follows that for every $g \in \overline{T}$

$$\phi_g(\overline{T}) \subseteq \overline{\phi_g(T)} \subseteq \overline{T},$$

as $\phi_g(T) \subseteq \overline{T}$, by the above consideration. So for any $g, h \in \overline{T}$, $g \cdot h = \phi_g(h) \in \overline{T}$. \square

Remark. Note that every preordering T on an \mathbb{R} -algebra E is a convex cone, so by (T2) and the Separation Theorem for convex sets (e.g. [5, II.39, Corollary 5]), an element x belongs to \overline{T} if and only if $L(x) \geq 0$ for every linear functional L on E such that $L(T) \geq 0$. It follows that for all elements $a, b \in E$ such that $a + \epsilon b \in T$ for every $\epsilon > 0$, we have that $a \in \overline{T}$.

From now on, we are mostly interested in localizations of the real polynomial algebra $\mathbb{R}[\underline{X}]$ in n variables $\underline{X} = (X_1, \dots, X_n)$. So for

a multiplicative set $D \subseteq \mathbb{R}[\underline{X}] \setminus \{0\}$ containing 1, we examine $E = D^{-1}\mathbb{R}[\underline{X}]$.

Remark. While the vector space dimension of $\mathbb{R}[\underline{X}]$ is countable, the dimension of E can be uncountable. For example, if D is the set of polynomials without real zeros, then the uncountable family $\left\{ \frac{1}{X_1^2 + c^2} \mid c \in \mathbb{R}^{\neq 0} \right\} \subseteq D^{-1}\mathbb{R}[\underline{X}]$ is linearly independent.

For a set $\mathcal{X} \subseteq \mathbb{R}^n$, an arbitrary function $\phi: \mathcal{X} \rightarrow \mathbb{R}$ and an element $R \in E = D^{-1}\mathbb{R}[\underline{X}]$, we say

$$R \succeq \phi \text{ on } \mathcal{X},$$

if there exist $f \in \mathbb{R}[\underline{X}]$ and $d \in D$ such that $R = \frac{f}{d}$ and $(f - d\phi)d \geq 0$ on \mathcal{X} . This just means that R has a representation $\frac{f}{d}$, $f \in \mathbb{R}[\underline{X}]$, $d \in D$, where the function $\frac{f}{d}$ is $\geq \phi$ pointwise on \mathcal{X} , wherever it is defined. The set

$$\text{Pos}^E(\mathcal{X}) = \{R \in E \mid R \succeq 0 \text{ on } \mathcal{X}\}$$

is a preordering in E . We also write

$$\text{Pos}(\mathcal{X}) = \{f \in \mathbb{R}[\underline{X}] \mid f \geq 0 \text{ on } \mathcal{X}\}$$

for the preordering of all polynomials nonnegative on \mathcal{X} in the usual sense. Note that $-1 \notin \text{Pos}(\mathcal{X})$ unless \mathcal{X} is empty. On the other hand, $-1 \in \text{Pos}^E(\mathcal{X})$ if $\mathcal{X} \subseteq \mathcal{Z}(d)$ for some $d \in D$. However, if $\mathcal{X} \setminus \mathcal{Z}(d)$ is dense in \mathcal{X} for every $d \in D$, such phenomena cannot occur.

Lemma 2. *For every element $R \in E = D^{-1}\mathbb{R}[\underline{X}]$ and every set $\mathcal{X} \subseteq \mathbb{R}^n$ such that $\mathcal{X} \setminus \mathcal{Z}(d)$ is dense in \mathcal{X} for every $d \in D$, the following are equivalent:*

- (1) $R \in \text{Pos}^E(\mathcal{X})$,
- (2) for every representation $R = \frac{f}{d}$ with $f \in \mathbb{R}[\underline{X}]$ and $d \in D$, we have that $fd \in \text{Pos}(\mathcal{X})$,
- (3) if we consider R as an element of $\mathbb{R}(\underline{X})$ and write $R = \frac{a}{b}$ where $a, b \in \mathbb{R}[\underline{X}]$, $b \neq 0$ and $\gcd(a, b) = 1$, then $ab \in \text{Pos}(\mathcal{X})$.

In particular, for all $f \in \mathbb{R}[\underline{X}]$, $\frac{f}{1} \in \text{Pos}^E(\mathcal{X})$ if and only if $f \in \text{Pos}(\mathcal{X})$.

Proof. Suppose that $R = \frac{f}{d} = \frac{a}{b}$ where $a, b \in \mathbb{R}[\underline{X}]$, $b \neq 0$, satisfy $\gcd(a, b) = 1$ and $f \in \mathbb{R}[\underline{X}]$, $d \in D$. Then there exists $u \in \mathbb{R}[\underline{X}] \setminus \{0\}$ such that $f = ua$ and $d = ub$. Clearly, $fd = abu^2 \in \text{Pos}(\mathcal{X})$ if and only if $ab \in \text{Pos}(\mathcal{X} \setminus \mathcal{Z}(bu))$. By the assumption on \mathcal{X} , this is equivalent to $ab \in \text{Pos}(\mathcal{X})$. So, (1), (2) and (3) are equivalent. The last claim is a special case of (3). \square

For a finite subset $S = \{g_1, \dots, g_m\}$ of $\mathbb{R}[\underline{X}]$ let

$$K_S = \{a \in \mathbb{R}^n \mid g_1(a) \geq 0, \dots, g_m(a) \geq 0\}$$

be the basic closed semialgebraic set defined by S . Let

$$T_S = \left\{ \sum_{\nu \in \{0,1\}^m} s_\nu g_1^{\nu_1} \cdots g_m^{\nu_m} \mid s_\nu \in \sum \mathbb{R}[\underline{X}]^2 \right\}$$

and

$$T_S^E = \left\{ \sum_{\nu \in \{0,1\}^m} s_\nu g_1^{\nu_1} \cdots g_m^{\nu_m} \mid s_\nu \in \sum E^2 \right\}$$

be the preorderings generated by S in $\mathbb{R}[\underline{X}]$ and $E = D^{-1}\mathbb{R}[\underline{X}]$, respectively. We always assume $K_S \neq \emptyset$, so $-1 \notin T_S$. If K_S is not contained in any $\mathcal{Z}(d)$, $d \in D$, then also $-1 \notin T_S^E$.

The following is our main result:

Theorem 1. *Let $D \subseteq \mathbb{R}[\underline{X}] \setminus \{0\}$ be a multiplicative set containing 1 and S a finite subset of $\mathbb{R}[\underline{X}]$. Suppose there is some $p \in D$ such that $p \geq 1$ on K_S and $kp - \sum_{i=1}^n X_i^2 \geq 0$ on K_S for some $k \geq 1$. Then*

$$\overline{T_S^E} = \text{Pos}^E(K_S \setminus \bigcup_{d \in D} \mathcal{Z}(d))$$

holds in $E = D^{-1}\mathbb{R}[\underline{X}]$.

The set $\mathcal{X} = K_S \setminus \bigcup_{d \in D} \mathcal{Z}(d)$ satisfies the assumptions of Lemma 2, which gives two characterizations of $\text{Pos}^E(\mathcal{X})$. Another characterization is that $R \in \text{Pos}^E(\mathcal{X})$ if and only if $\chi(R) \geq 0$ for every unital \mathbb{R} -algebra homomorphism $\chi: E \rightarrow \mathbb{R}$ such that $\chi(S) \geq 0$.

We will give the proof of Theorem 1 in Section 3. In Section 4, we will show that Theorem 1 implies the solution of the real multidimensional rational K -moment problem. In Section 5 we will prove a variant of Theorem 1 for the topology of finitely open sets.

Note that in case K_S is compact, we can always take $p = 1$ in the theorem. In the noncompact case, the polynomial $p = 1 + \sum_{i=1}^n X_i^2$ can always be used in the application of Theorem 1, as long as it belongs to the multiplicative set D . We record an easy corollary of Theorem 1, in which this is the case. Therefore let $\mathbb{R}(\underline{X}; \mathcal{P})$ be the algebra of real rational functions with (real) poles only in a given set $\mathcal{P} \subseteq \mathbb{R}^n$. It is the localization of the polynomial algebra with respect to the set D of nonzero polynomials with real zeros only in \mathcal{P} . The set of zeros of elements from D equals \mathcal{P} . The polynomial $p = 1 + \sum_{i=1}^n X_i^2$ belongs to D .

Corollary 1. *In $E = \mathbb{R}(\underline{X}; \mathcal{P}) = D^{-1}\mathbb{R}[\underline{X}]$ we have for arbitrary finite sets $S \subseteq \mathbb{R}[\underline{X}]$*

$$\overline{T_S^E} = \text{Pos}^E(K_S \setminus \mathcal{P}).$$

In particular

$$\overline{\sum \mathbb{R}(\underline{X}; \mathcal{P})^2} = \text{Pos}^E(\mathbb{R}^n \setminus \mathcal{P}).$$

If $\mathcal{P} = \mathbb{R}^n$, then $\mathbb{R}(\underline{X}; \mathcal{P}) = \mathbb{R}(\underline{X})$ and $\text{Pos}^E(\mathbb{R}^n \setminus \mathcal{P}) = \mathbb{R}(\underline{X})$, so

$$\overline{\sum \mathbb{R}(\underline{X})^2} = \mathbb{R}(\underline{X}).$$

In particular, there is no positive linear functional on $\mathbb{R}(\underline{X})$.

The last claim of Corollary 1 also follows from [14, 9.7.29].

Remark. Let L/\mathbb{R} be a proper field extension. We claim that there is no nontrivial positive linear functional on L . If L is not real, then every element is a sum of squares, so there is clearly no nontrivial positive linear functional. Otherwise, every element a of $L \setminus \mathbb{R}$ is transcendental over \mathbb{R} . So $\mathbb{R}(a)$ is isomorphic to $\mathbb{R}(X)$, and by Corollary 1 there is no nontrivial positive linear functional on $\mathbb{R}(a)$.

Remark. When K_S is compact and disjoint from $\bigcup_{d \in D} \mathcal{Z}(d)$, then we can prove Theorem 1 by the usual analytic trick. Namely, in this case Wörmann's trick implies that T_S^E is archimedean. (Since Wörmann's trick works only for finitely generated \mathbb{R} -algebras, we apply it first to algebras $\mathbb{R}[\underline{X}]_d$ and then use that $E = D^{-1}\mathbb{R}[\underline{X}]$ is equal to $\bigcup_{d \in D} \mathbb{R}[\underline{X}]_d$.) Then the Kadison-Dubois Representation Theorem implies that the closure of T_S^E is equal to $\text{Pos}^E(K_S)$. We refer the reader to [11] for the statement of the Wörmann's trick and the Representation Theorem. We will not pursue this idea further.

3. THE PROOF OF THEOREM 1

The easy inclusion $\overline{T_S^E} \subseteq \text{Pos}^E(K_S \setminus \bigcup_{d \in D} \mathcal{Z}(d))$ follows from the obvious fact that evaluation in a point from $K_S \setminus \bigcup_{d \in D} \mathcal{Z}(d)$ defines a linear functional L on E such that $L(T_S^E) \geq 0$.

The proof of the difficult inclusion $\overline{T_S^E} \supseteq \text{Pos}^E(K_S \setminus \bigcup_{d \in D} \mathcal{Z}(d))$ will be split into several lemmas.

Lemma 3. *Let $D \subseteq \mathbb{R}[\underline{X}] \setminus \{0\}$ be a multiplicative set containing 1, K a compact set contained in $\mathcal{P} = \bigcup_{d \in D} \mathcal{Z}(d)$, and $c \in \mathbb{R}^{>0}$. Then every neighbourhood of zero in $D^{-1}\mathbb{R}[\underline{X}]$ (in the finest locally convex topology) contains an element R such that $R \succeq c \cdot \chi_K$ globally, where χ_K is the characteristic function of K .*

Proof. Let N be a neighborhood of zero in $D^{-1}\mathbb{R}[X]$, without loss of generality assume that N is convex and absorbent. So for any $f \in D^{-1}\mathbb{R}[X]$ there is a number $\delta(f) > 0$, such that $\lambda f \in N$ for all $\lambda \in [-\delta(f), \delta(f)]$.

For every $a \in \bigcup_{d \in D} \mathcal{Z}(d)$ we have $d_a(a) = 0$ for some $d_a \in D$, so that $d_a(x) = (\nabla d_a)(a) \cdot (x - a) + o(\|x - a\|)$. Write $c_a = \|(\nabla d_a)(a)\| + 1$ and pick $\eta_a > 0$ such that

$$(1) \quad |d_a(x)| \leq c_a \|x - a\| \text{ on } B(a, \eta_a).$$

Let W be a cube containing K and let B_n be the unit ball in \mathbb{R}^n . Define

$$(2) \quad \lambda := \frac{1}{6} \left(\frac{\text{vol}(B_n)}{c \text{vol}(W)} \right)^{\frac{1}{n}}$$

and

$$(3) \quad r_a := \min\{\lambda c_a^{-2} \delta(d_a^{-2n})^{\frac{1}{n}}, \text{vol}(W)^{\frac{1}{n}}, \eta_a\}.$$

So $r_a > 0$ and

$$K \subseteq \bigcup_{a \in K} B(a, \frac{r_a}{3}),$$

where $B(a, \frac{r_a}{3})$ denotes the open ball of radius $\frac{r_a}{3}$ around a . By Wiener's Covering Lemma, [10, Lemma 4.1.1], there are $a_1, \dots, a_t \in K$ such that

$$K \subseteq \bigcup_{i=1}^t B(a_i, r_{a_i})$$

and the $B(a_i, \frac{r_{a_i}}{3})$ are pairwise disjoint.

For every ball B with center in W and with radius less or equal than half the side length of W we have

$$\text{vol}(B \cap W) \geq \frac{1}{2^n} \text{vol}(B),$$

since $\text{vol}(B \cap W)$ is minimal if the center of B lies in one of the corners of W . In particular, this is true for $B = B(a_i, \frac{r_{a_i}}{3})$, $i = 1, \dots, t$, by (3). Since $B(a_i, \frac{r_{a_i}}{3})$ are pairwise disjoint, it follows that

$$\sum_{i=1}^t \frac{1}{2^n} \text{vol}\left(B(a_i, \frac{r_{a_i}}{3})\right) \leq \text{vol}(W).$$

Combining this with the volume formula

$$\text{vol}\left(B(a_i, \frac{r_{a_i}}{3})\right) = \left(\frac{r_{a_i}}{3}\right)^n \text{vol}(B_n),$$

we get

$$(4) \quad \sum_{i=1}^t r_{a_i}^n \leq \frac{6^n \operatorname{vol}(W)}{\operatorname{vol}(B_n)} = \frac{1}{\lambda^n c},$$

where the last equality follows from (2). Now define

$$R := \frac{1}{\lambda^n \sum_{i=1}^t r_{a_i}^n} \cdot \sum_{i=1}^t \frac{r_{a_i}^{2n} c_{a_i}^{2n}}{d_{a_i}^{2n}} = \sum_{i=1}^t \frac{r_{a_i}^n}{\sum_{j=1}^t r_{a_j}^n} \cdot \frac{r_{a_i}^n c_{a_i}^{2n}}{\lambda^n d_{a_i}^{2n}}.$$

Clearly $R \in D^{-1}\mathbb{R}[\underline{X}]$ and, as $\frac{r_{a_i}^n c_{a_i}^{2n}}{\lambda^n} \leq \delta(d_{a_i}^{-2n})$ by (3), each $\frac{r_{a_i}^n c_{a_i}^{2n}}{\lambda^n d_{a_i}^{2n}}$ lies in N . So R , as a convex combination of such elements, also lies in N . By (1) and (3), $\frac{r_{a_i}^{2n} c_{a_i}^{2n}}{d_{a_i}^{2n}} \succeq \chi_{B(a_i, r_{a_i})}$ globally, so one checks that

$$\sum_{i=1}^t \frac{r_{a_i}^{2n} c_{a_i}^{2n}}{d_{a_i}^{2n}} \succeq \chi_K.$$

Therefore, $R \succeq c \cdot \chi_K$ by (4). \square

Lemma 4. *Let $D \subseteq \mathbb{R}[\underline{X}] \setminus \{0\}$ be a multiplicative set containing 1 and $S \subseteq \mathbb{R}[\underline{X}]$ finite. Assume there is some $p \in D$ such that $p \geq 1$ on K_S , and $kp - \sum_{i=1}^n X_i^2 \geq 0$ on K_S for some $k \geq 1$. Let K be a compact subset of $\mathcal{P} = \bigcup_{d \in D} \mathcal{Z}(d)$. Then every $f \in \mathbb{R}[\underline{X}]$ which is nonnegative on $K_S \setminus K$ belongs to $\overline{T_S^E}$ in $E = D^{-1}\mathbb{R}[\underline{X}]$.*

Proof. Take any neighborhood of zero N in E . Let m be the minimum of f on $K_S \cap K$. By Lemma 3, there exists $R \in N$ such that $R \succeq -m \cdot \chi_K$. So we find a representation $f + R = \frac{u}{v}$ with $uv \geq 0$ on K_S . By [22, Theorem 5.1]¹, the polynomial uv belongs to the closure of the preordering generated by S in $\mathbb{R}[\underline{X}, \frac{1}{p}]$, so also to the closure of T_S^E . So $f + R = (\frac{1}{v})^2 uv \in (f + N) \cap \overline{T_S^E}$. So $f \in \overline{T_S^E}$, as N was arbitrary. \square

In the next Lemma we get rid of the compactness assumption:

Lemma 5. *Lemma 4 also holds when $K = \mathcal{P}$.*

Proof. For every $\epsilon > 0$, $f + \epsilon p^{\deg(f)+1}$ is nonnegative on $K_S \setminus B_\epsilon$ for a closed ball B_ϵ around 0. Define

$$K_\epsilon := \{a \in K_S \cap B_\epsilon \mid f(x) \leq -\epsilon\}.$$

¹Note that our assumption on p implies that condition (*) of [22, Theorem 5.1] is satisfied. If p satisfies, instead of $p \geq 1$ on K_S , the stronger assumption $p - 1 \in T_S$ - e.g. when $p = 1 + \sum_{i=1}^n X_i^2$ - then we can use [11, Theorem 6.2.3] (or [12, Corollary 3.2]), which is a slightly weaker version of [22, Theorem 5.1].

K_ϵ is a compact subset of $\mathcal{P} = \bigcup_{d \in D} \mathcal{Z}(d)$ and $f + \epsilon + \epsilon p^{\deg(f)+1}$ is nonnegative on $K_S \setminus K_\epsilon$. By Lemma 4, $f + \epsilon + \epsilon p^{\deg(f)+1}$ belongs to the closure of T_S^E in $E = D^{-1}\mathbb{R}[\underline{X}]$. As this is true for all $\epsilon > 0$, f belongs to $\overline{T_S^E}$. \square

Now we give the

Proof of Theorem 1. Take $\frac{f}{d} \in E$ with $fd \in \text{Pos}(K_S \setminus \mathcal{P})$. Then apply Lemma 5 to obtain $fd \in \overline{T_S^E}$ and multiply with $(\frac{1}{d})^2$. \square

4. RATIONAL MOMENT PROBLEMS

The aim of this section is to prove the following existence result for the multidimensional rational K -moment problem, which in the one-dimensional compact case extends [8, Theorem 5].

Theorem 2. *Let $D \subseteq \mathbb{R}[\underline{X}] \setminus \{0\}$ be a multiplicative set containing 1 and let $S \subseteq \mathbb{R}[\underline{X}]$ be finite. Assume there is some $p \in D$ such that $p \geq 1$ on K_S , and $kp - \sum_{i=1}^n X_i^2 \geq 0$ on K_S for some $k \geq 1$.*

Then, for every linear functional L on $E = D^{-1}\mathbb{R}[\underline{X}]$ such that $L(T_S^E) \geq 0$, there exists a measure μ on $\overline{K_S \setminus \bigcup_{d \in D} \mathcal{Z}(d)}$ such that

$$L\left(\frac{f}{d}\right) = \int \frac{f}{d} d\mu$$

for every $\frac{f}{d} \in D^{-1}\mathbb{R}[\underline{X}]$.

If K_S has empty intersection with $\mathcal{P} = \bigcup_{d \in D} \mathcal{Z}(d)$, then Theorem 2 follows from our Theorem 1 and [11, Theorem 3.2.2]. To prove the general case, we need the following generalization of [11, Theorem 3.2.2] (applied to $A = \mathbb{R}[\underline{X}]$, $\mathcal{X} = \overline{K_S \setminus \bigcup_{d \in D} \mathcal{Z}(d)}$, $\hat{a} = a|_{\mathcal{X}}$ and $q = \sum X_i^2$.)

Theorem 3 (Rational Haviland's Theorem). *Let A be a unital \mathbb{R} algebra, D a multiplicative subset of A containing 1, \mathcal{X} a nonempty Hausdorff topological space and $\hat{\cdot} : A \rightarrow \mathcal{C}(\mathcal{X}, \mathbb{R})$ an \mathbb{R} -algebra homomorphism. Suppose that:*

- (1) *there exists $q \in A$ such that $\hat{q} \geq 0$ on \mathcal{X} and, for each $k \geq 1$, the set $\mathcal{X}_k = \{x \in \mathcal{X} \mid \hat{q}(x) \leq k\}$ is compact,*
- (2) *for every $d \in D$, the zero set $\mathcal{Z}(\hat{d})$ of \hat{d} in \mathcal{X} has empty interior,*
- (3) *every open subset of \mathcal{X} is σ -compact (= a countable union of compact sets).*

Then, for any linear functional $L : D^{-1}A \rightarrow \mathbb{R}$, satisfying

$$\forall a \in A, \forall d \in D, \hat{a}\hat{d} \geq 0 \text{ on } \mathcal{X} \Rightarrow L(a/d) \geq 0,$$

there exists a Borel measure μ on \mathcal{X} such that

$$\forall a \in A, \forall d \in D, L(a/d) = \int_{\mathcal{X}} \hat{a}/\hat{d} \, d\mu.$$

The idea is to follow the proof of [11, Theorem 3.2.2] (or [13, Theorem 3.1]) and use [24, Theorem 3.2] instead of Riesz's Theorem. We start with a few remarks:

R1. Assumption (1) implies that \mathcal{X} is locally compact.

R2. For every $d \in D$, the following assertions are equivalent:

- (a) $\mathcal{X} \setminus \mathcal{Z}(\hat{d})$ is dense in \mathcal{X} ,
- (b) $\mathcal{Z}(\hat{d})$ has empty interior,
- (c) for every $f \in \mathcal{C}(\mathcal{X}, \mathbb{R})$, if $f\hat{d}^2 \geq 0$ on \mathcal{X} then $f \geq 0$ on \mathcal{X} ,
- (d) for every $f \in \mathcal{C}(\mathcal{X}, \mathbb{R})$, if $f\hat{d} \equiv 0$ on \mathcal{X} then $f \equiv 0$ on \mathcal{X} .

Every locally compact Hausdorff space is completely regular, which means that points can be separated from closed sets by continuous functions (Urysohn's Lemma, see [19, 2.12]). So we have that (d) \Rightarrow (b). Implications (a) \Leftrightarrow (b) \Rightarrow (c) \Rightarrow (d) are clear.

R3. Assumption (2) implies that $0 \notin \hat{D}$, hence $0 \notin D$. Therefore, $D^{-1}A$ and $\hat{D}^{-1}\mathcal{C}(\mathcal{X}, \mathbb{R})$ are nontrivial \mathbb{R} -algebras.

R4. The homomorphism $\hat{\cdot}: A \rightarrow \mathcal{C}(\mathcal{X}, \mathbb{R})$ extends uniquely to a homomorphism $\hat{\cdot}: D^{-1}A \rightarrow \widehat{\hat{D}^{-1}\mathcal{C}(\mathcal{X}, \mathbb{R})}$. Its image is $\widehat{D^{-1}A} = \hat{D}^{-1}\hat{A}$.

R5. $\bar{L}: \hat{D}^{-1}\hat{A} \rightarrow \mathbb{R}$ defined by $\bar{L}(\hat{a}/\hat{d}) = L(a/d)$ is well-defined. This follows from the positivity assumption on L . (See Claim 1 in the proof of [11, Theorem 3.2.2].)

R6. We can assume that $\hat{d} \geq 0$ on \mathcal{X} for every $d \in D$. Namely, every element $f/\hat{d} \in \hat{D}^{-1}\mathcal{C}(\mathcal{X}, \mathbb{R})$ can be written as $f/\hat{d} = f\hat{d}/\hat{d}^2$ and $\hat{d}^2 \geq 0$ on \mathcal{X} . The positivity assumption on L then simplifies to: $\forall a \in A, \forall d \in D, \hat{a} \geq 0$ on $\mathcal{X} \Rightarrow L(a/d) \geq 0$.

R7. Let $B = \hat{D}^{-1}\mathcal{C}'(\mathcal{X}, \mathbb{R})$, where $\mathcal{C}'(\mathcal{X}, \mathbb{R}) = \{f \in \mathcal{C}(\mathcal{X}, \mathbb{R}) \mid \exists a \in A: |f| \leq \hat{a} \text{ on } \mathcal{X}\}$. Note that B is a subalgebra of $\hat{D}^{-1}\mathcal{C}(\mathcal{X}, \mathbb{R})$ and $B = \{f/\hat{d} \in \hat{D}^{-1}\mathcal{C}(\mathcal{X}, \mathbb{R}) \mid \exists a \in A, s \in D: |f|\hat{s} \leq \hat{a}\hat{d} \text{ on } \mathcal{X}\}$. Namely, if $|f|\hat{s} \leq \hat{a}\hat{d}$ on \mathcal{X} , then $f/\hat{d} = f\hat{s}/\hat{d}\hat{s}$ and $f\hat{s} \in \mathcal{C}'(\mathcal{X}, \mathbb{R})$. The other inclusion is clear.

R8. The algebra B obviously contains both $\hat{D}^{-1}\hat{A}$ and $\hat{D}^{-1}\mathcal{C}_c(\mathcal{X}, \mathbb{R})$. Here, $\mathcal{C}_c(\mathcal{X}, \mathbb{R})$ denotes the algebra of continuous functions with compact support.

Lemma 6. \bar{L} extends to a linear functional \bar{L} on $B = \hat{D}^{-1}\mathcal{C}'(\mathcal{X}, \mathbb{R})$ such that $\bar{L}(f/\hat{d}) \geq 0$ for every $d \in D$ and every nonnegative $f \in \mathcal{C}'(\mathcal{X}, \mathbb{R})$.

Proof. The proof is similar as the proof of Claim 2 in the proof of [11, Theorem 3.2.2].) This result is also a special case of a more general extension theorem for positive functionals, see [2, Theorem 2.6.2, p. 69]. \square

Lemma 7 (Rational Riesz's Theorem). *For every linear functional $\bar{L}: \hat{D}^{-1}\mathcal{C}_c(\mathcal{X}, \mathbb{R}) \rightarrow \mathbb{R}$ such that $\bar{L}(f/\hat{d}) \geq 0$ for every nonnegative $f \in \mathcal{C}_c(\mathcal{X}, \mathbb{R})$ there exists a unique Borel measure μ on \mathcal{X} such that $\bar{L}(f/\hat{d}) = \int_{\mathcal{X}} f/\hat{d} d\mu$ for every $f/\hat{d} \in \hat{D}^{-1}\mathcal{C}_c(\mathcal{X}, \mathbb{R})$.*

Proof. (Based on the proof of [24, Theorem 3.2].) For every $s \in D$ define a functional $\bar{L}_s: \mathcal{C}_c(\mathcal{X}, \mathbb{R}) \rightarrow \mathbb{R}$ by $\bar{L}_s(f) = \bar{L}(f/\hat{s})$. Since $\bar{L}_s(f) \geq 0$ for every $f \geq 0$, there exists by Riesz's Theorem, see [19, Theorem 2.14], a unique regular Borel measure μ_s on \mathcal{X} such that $\bar{L}_s(f) = \int_{\mathcal{X}} f d\mu_s$ for every $f \in \mathcal{C}_c(\mathcal{X}, \mathbb{R})$. Write $\mu = \mu_1$ and note that

$$\int_{\mathcal{X}} f d\mu = \int_{\mathcal{X}} f\hat{s} d\mu_s$$

for every $f \in \mathcal{C}_c(\mathcal{X}, \mathbb{R})$. By [19, Theorem 1.29], $\hat{s} d\mu_s$ is a Borel measure on \mathcal{X} . To prove that $\hat{s} d\mu_s$ is regular, we apply [19, Theorem 2.18]. (We need here our assumption (3) and continuity of \hat{s} .) By the uniqueness part of Riesz's Theorem, $\mu = \hat{s} d\mu_s$.

Now, we will prove that $\mu_s(\mathcal{Z}(\hat{s})) = 0$. By a version of Urysohn's Lemma (see [19, Theorem 2.12]), there exists for each integer k a function $u_k \in \mathcal{C}_c(\mathcal{X}, \mathbb{R})$ such that $0 \leq u_k \leq 1$ on \mathcal{X} and $u_k = 1$ on $\mathcal{Z}(\hat{s}) \cap \mathcal{X}_k$. For all integers i, k , the function $f_{k,i} = u_k \min\{1, 1/i\hat{s}\}$ belongs to $\mathcal{C}_c(\mathcal{X}, \mathbb{R})$, $0 \leq f_{k,i} \leq 1$ on \mathcal{X} and $f_{k,i} = 1$ on $\mathcal{Z}(\hat{s}) \cap \mathcal{X}_k$. It follows that

$$\begin{aligned} \mu_s(\mathcal{Z}(\hat{s}) \cap \mathcal{X}_k) &\leq \int_{\mathcal{X}} f_{k,i} d\mu_s = \\ &= \bar{L}_s(f_{k,i}) = \bar{L}(f_{k,i}/\hat{s}) \leq (1/i)\bar{L}(u_k/\hat{s}^2). \end{aligned}$$

Sending $i \rightarrow \infty$, we get $\mu_s(\mathcal{Z}(\hat{s}) \cap \mathcal{X}_k) = 0$ for every k , hence $\mu_s(\mathcal{Z}(\hat{s})) = 0$. In particular, $\mu(\mathcal{Z}(\hat{s})) = 0$ (which is also a consequence of $\mu = \hat{s} d\mu_s$). Finally, for every $f \in \mathcal{C}_c(\mathcal{X}, \mathbb{R})$ and every $s \in D$ we have that

$$\int_{\mathcal{X}} f/\hat{s} d\mu = \int_{\mathcal{X} \setminus \mathcal{Z}(\hat{s})} f/\hat{s} d\mu = \int_{\mathcal{X} \setminus \mathcal{Z}(\hat{s})} (f/\hat{s})\hat{s} d\mu_s =$$

$$= \int_{\mathcal{X} \setminus \mathcal{Z}(\hat{s})} f \, d\mu_s = \int_{\mathcal{X}} f \, d\mu_s = \bar{L}_s(f) = \bar{L}(f/\hat{s}).$$

□

Lemma 8. *For every nonnegative function $f \in \mathcal{C}'(\mathcal{X}, \mathbb{R})$, there exists a monotonically increasing sequence of nonnegative functions $f_i \in \mathcal{C}_c(\mathcal{X}, \mathbb{R})$ such that $\frac{(f+\hat{q})^2}{i} \geq f - f_i \geq 0$ on \mathcal{X} for every i .*

This is Claim 3 in the proof of [11, Theorem 3.2.2]. No changes are needed.

Proof of the Theorem. Let \bar{L} be a linear functional on $\hat{D}^{-1}\hat{A}$ such that $\bar{L}(f/\hat{d}) \geq 0$ for every nonnegative $f \in \hat{A}$ and every $d \in D$. By Lemma 6, we can extend \bar{L} to a linear functional \bar{L} on $\hat{D}^{-1}\mathcal{C}'(\mathcal{X}, \mathbb{R})$ such that $\bar{L}(f/\hat{d}) \geq 0$ for every nonnegative $f \in \mathcal{C}'(\mathcal{X}, \mathbb{R})$. Lemma 7 gives us a Borel measure μ on \mathcal{X} such that $\bar{L}(f/\hat{d}) = \int_{\mathcal{X}} f/\hat{d} \, d\mu$ for every $f \in \mathcal{C}_c(\mathcal{X}, \mathbb{R})$. Lemma 8 implies that $\bar{L}(f/\hat{d}) = \int_{\mathcal{X}} f/\hat{d} \, d\mu$ for every $f \in \mathcal{C}'(\mathcal{X}, \mathbb{R})$. Namely,

$$\int_{\mathcal{X}} f/\hat{d} \, d\mu = \lim_{i \rightarrow \infty} \int_{\mathcal{X}} f_i/\hat{d} \, d\mu = \lim_{i \rightarrow \infty} \bar{L}(f_i/\hat{d}) = \bar{L}(f/\hat{d}),$$

by the Monotone Convergence Theorem, the inequality $\frac{(f+\hat{q})^2}{i} \geq f - f_i \geq 0$ and the positivity of \bar{L} . In particular $L(a/d) = \bar{L}(\hat{a}/\hat{d}) = \int_{\mathcal{X}} \hat{a}/\hat{d} \, d\mu$ for every $a \in A$ and $d \in D$. □

5. PREORDERINGS AND THE TOPOLOGY OF FINITELY OPEN SETS

Definition. Let E be an \mathbb{R} -vector space and O a subset of E . We say that O is open in the *topology of finitely open sets* if for every finite-dimensional vector subspace U of E , $O \cap U$ is open in the usual topology of U . Write \mathcal{T}_{fin} for the set of all open sets.

We recall the following results from [3].

- (F1) If E has countable dimension, then the topology \mathcal{T}_{fin} is equal to the topology \mathcal{T}_ω . If E does not have countable dimension, then the topology \mathcal{T}_{fin} is strictly finer (=has more open sets) than the topology \mathcal{T}_ω . In particular, \mathcal{T}_{fin} is not a locally convex topology in this case.
- (F2) If E does not have countable dimension, then the addition is not continuous in the topology \mathcal{T}_{fin} .

(F3) Let E be an \mathbb{R} -vector space and \mathcal{D} the set of all countable-dimensional subspaces of E . Then, for every subset A of E

$$\overline{A} = \bigcup_{D \in \mathcal{D}} \overline{A \cap D}.$$

The following should be well-known but we were unable to find a reference:

(F4) Any linear mapping between any two vector spaces both having the topology of finitely open sets is continuous. In particular, every functional on every vector space with the finitely open topology is continuous.

Proof. Let $L: (E, \mathcal{T}_{\text{fin}}) \rightarrow (E', \mathcal{T}'_{\text{fin}})$ be a linear mapping and $O \in \mathcal{T}'_{\text{fin}}$. Pick any finite-dimensional subspace V of E . Clearly, the subspace $L(V)$ of E' is also finite-dimensional. By the definition of $\mathcal{T}'_{\text{fin}}$, $O \cap L(V)$ is open in the usual topology of $L(V)$. Every linear mapping between two finite-dimensional vector spaces is continuous in the usual topologies. In particular $L|_V: V \rightarrow L(V)$ is continuous in the usual topologies of V and $L(V)$. It follows, that the set $(L|_V)^{-1}(O \cap L(V))$ is open in the usual topology of V . Note that $(L|_V)^{-1}(O \cap L(V)) = L^{-1}(O) \cap V$, hence $L^{-1}(O) \cap V$ is open in the usual topology of V . Since V was arbitrary, it follows by the definition of \mathcal{T}_{fin} that $L^{-1}(O)$ is open. Since O was arbitrary, L is indeed continuous. \square

Lemma 9. *For a preordering T on a commutative \mathbb{R} -algebra E , its closure in \mathcal{T}_{fin} is also a preordering.*

Proof. For any $f_1, f_2 \in \overline{T}$, there exist, by (F3), countable-dimensional subspaces D_1, D_2 of E such that $f_i \in \overline{T \cap D_i}$ for $i = 1, 2$. The subspace $D = D_1 + D_2$ is also countable-dimensional and $\overline{T \cap D_i} \subseteq \overline{T \cap D}$ for $i = 1, 2$. It follows that $f_1, f_2 \in \overline{T \cap D}$. Since D is countable-dimensional, $\overline{T \cap D}$ is closed under addition, hence $f_1 + f_2 \in \overline{T \cap D} \subseteq \overline{T}$. The proof that \overline{T} is closed under multiplication follows again from the fact, that $u \mapsto ru$ is continuous for all $r \in E$. (See the proof of Lemma 1). \square

Lemma 10. *Let $D \subseteq \mathbb{R}[\underline{X}] \setminus \{0\}$ be a multiplicative set containing 1. Then a preordering T in $E = D^{-1}\mathbb{R}[\underline{X}]$ is closed if and only if $T \cap \mathbb{R}[\underline{X}]$ is closed in $\mathbb{R}[\underline{X}]$.*

Proof. One direction is obvious. So assume that $T \cap \mathbb{R}[\underline{X}]$ is closed in $\mathbb{R}[\underline{X}]$. Let V be a finite-dimensional subspace of E . We claim that $V \cap T$ is closed. Pick a nonzero square $d \in D$ such that $dV \subseteq \mathbb{R}[\underline{X}]$. Then

$$d(V \cap T) = dV \cap dT = dV \cap T.$$

The first equality follows from the fact that the mapping $u \mapsto du$ is bijective on E and the second from $T = dT$ (d is an invertible square in E). Since $T \cap \mathbb{R}[\underline{X}]$ is closed and $dV \subseteq \mathbb{R}[\underline{X}]$, $d(V \cap T)$ is closed in dV . As the mapping $u \mapsto du$ is a homeomorphism by (F4), it follows that the set $V \cap T$ is closed in V . As V was arbitrary, T is closed in E . \square

The next Lemmas are needed in the proof of the main theorem of this section. See [4, Chapter 2] for general notions such as semialgebraic set, semialgebraic homeomorphism, semialgebraic dimension etc.

Lemma 11. *Let $K \subseteq \mathbb{R}^n$ be a semialgebraic set. Then there are finitely many semialgebraic subsets C_1, \dots, C_s of K , such that*

- (i) $K = C_1 \cup \dots \cup C_s$,
- (ii) each C_i is semialgebraically homeomorphic to some \mathbb{R}^{d_i} ,
- (iii) each C_i has irreducible Zariski closure.

Proof. The proof is by induction on the semialgebraic dimension of K . If $\dim(K) = 0$, the result is clear, as K is a finite union of points. So assume $\dim(K) = m$ and the result is true for all semialgebraic sets of smaller dimension. Let V be the Zariski closure of K and V_1, \dots, V_r its irreducible components. We consider with out loss of generality only $K_1 := K \cap V_1$. The usual semialgebraic cell decomposition allows us to write

$$K_1 = \tilde{C}_1 \cup \dots \cup \tilde{C}_k,$$

where each \tilde{C}_j is a semialgebraic set and semialgebraically homeomorphic to some \mathbb{R}^{d_j} . Now if a \tilde{C}_j has the same dimension as V_1 , its Zariski closure equals V_1 , due to the irreducibility of V_1 . So we use all these C_j for our desired decomposition of K . The union of all lower dimensional C_j is a semialgebraic set of strictly smaller dimension than m , so we can apply the induction hypothesis to it. This yields the result. \square

Lemma 12. *For every semialgebraic set K in \mathbb{R}^n and every nonempty multiplicative subset D of $\mathbb{R}[\underline{X}]$, there exists an element $d_K \in D$ such that $\overline{K \setminus \mathcal{Z}(d_K)} \subseteq \overline{K \setminus \mathcal{Z}(d)}$ for every $d \in D$.*

Proof. Write $K = C_1 \cup \dots \cup C_s$ with the properties from Lemma 11. For each $i \in \{1, \dots, s\}$ and $d \in D$ there are two possibilities. First, $\dim(C_i \cap \mathcal{Z}(d)) < \dim(C_i)$. Since semialgebraic homeomorphisms respect dimensions ([4, Theorem 2.8.8]), we see that $C_i \cap \mathcal{Z}(d)$ has (relative) empty interior in C_i , so $\overline{C_i \setminus \mathcal{Z}(d)} = \overline{C_i}$. The second possibility is $\dim(C_i \cap \mathcal{Z}(d)) = \dim(C_i)$. Since the Zariski closure of C_i is irreducible, it equals the Zariski closure of $C_i \cap \mathcal{Z}(d)$. But this means d must vanish

on the whole of C_i . So $\overline{C_i \setminus \mathcal{Z}(d)} = \emptyset$. So we have proved that for every $d \in D$,

$$\overline{K \setminus \mathcal{Z}(d)} = \bigcup_{i=1}^s \overline{C_i \setminus \mathcal{Z}(d)} = \bigcup_{i \in I(d)} \overline{C_i},$$

where $I(d) = \{i \mid \dim(C_i \cap \mathcal{Z}(d)) < \dim(C_i)\}$. There exist finitely many $d_1, \dots, d_t \in D$ such that $\bigcap_{d \in D} I(d) = \bigcap_{j=1}^t I(d_j)$. Set $d_K := \prod_{j=1}^t d_j$ and note that $I(d_K) = \bigcap_{j=1}^t I(d_j)$. It follows that for every $d \in D$,

$$\overline{K \setminus \mathcal{Z}(d_K)} = \bigcup_{i \in I(d_K)} \overline{C_i} \subseteq \bigcup_{i \in I(d)} \overline{C_i} = \overline{K \setminus \mathcal{Z}(d)}.$$

□

Lemma 13. *For any semialgebraic set $K \subseteq \mathbb{R}^n$, the preordering*

$$\text{Pos}^E(K) = \{R \in D^{-1}\mathbb{R}[\underline{X}] \mid R \succeq 0 \text{ on } K\}$$

is closed in \mathcal{T}_{fin} .

Proof. Lemma 12 implies that the set $\mathcal{X} = \overline{K \setminus \mathcal{Z}(d_K)}$ satisfies the assumptions of Lemma 2, hence $\text{Pos}^E(\mathcal{X}) \cap \mathbb{R}[\underline{X}] = \text{Pos}(\mathcal{X})$. The set $\text{Pos}(\mathcal{X})$ is closed in $\mathbb{R}[\underline{X}]$, as evaluations in points are linear and therefore continuous. So $\text{Pos}^E(\mathcal{X})$ is closed by Lemma 10.

We also have that $\text{Pos}^E(\mathcal{X}) = \text{Pos}^E(K)$. Namely, if $R = \frac{f}{d}$ with $f \in \mathbb{R}[\underline{X}]$, and $d \in D$ such that $fd \in \text{Pos}(K \setminus \mathcal{Z}(d_K))$, then also $R = \frac{fd_K}{dd_K}$ with $(fd_K)(dd_K) \in \text{Pos}(K)$. The opposite inclusion is trivial. □

The following main theorem of this section is an analogue of Theorem 1 above.

Theorem 4. *Let $D \subseteq \mathbb{R}[\underline{X}] \setminus \{0\}$ be a multiplicative set containing 1 and $S \subseteq \mathbb{R}[\underline{X}]$ finite. Suppose there is $p \in D$ with $p \geq 1$ on K_S and $kp - \sum_{i=1}^n X_i^2 \geq 0$ on K_S for some $k \geq 1$. Then in the topology \mathcal{T}_{fin} on $D^{-1}\mathbb{R}[\underline{X}]$ we have*

$$\overline{T_S^E} = \text{Pos}^E(K_S).$$

Proof. First take $R \in \text{Pos}^E(K_S)$, so $R = \frac{f}{d}$ for some $f \in \mathbb{R}[\underline{X}]$, $d \in D$, with $fd \geq 0$ on K_S . We apply [22, Theorem 5.1] and find fd lying in the closure of T_S^E in a finite dimensional subspace of $D^{-1}\mathbb{R}[\underline{X}]$. So fd and therefore R belong to $\overline{T_S^E}$.

Since $T_S^E \subseteq \text{Pos}^E(K_S) \subseteq \overline{T_S^E}$, it remains to show that $\text{Pos}^E(K_S)$ is closed in \mathcal{T}_{fin} . This is Lemma 13. □

Remark. We can also deduce Theorem 4 from Theorem 1 and Lemma 13. Namely,

$$\bigcup_{d \in D} \text{Pos}^{\mathbb{R}[\underline{X}]_d}(K_S) \subseteq \bigcup_{d \in D} \text{Pos}^{\mathbb{R}[\underline{X}]_{pd}}(K_S) \subseteq \bigcup_{d \in D} \overline{T_S^{\mathbb{R}[\underline{X}]_{pd}}} \subseteq \bigcup_{d \in D} \overline{T_S^{\mathbb{R}[\underline{X}]_d}},$$

by application of Theorem 1 to the algebra $\mathbb{R}[\underline{X}]_{pd}$; we use that \mathcal{T}_ω and \mathcal{T}_{fin} coincide in the countable dimensional case. Now, using Lemma 13,

$$\overline{T_S^E} \subseteq \text{Pos}^E(K_S) \subseteq \bigcup_{d \in D} \text{Pos}^{\mathbb{R}[\underline{X}]_d}(K_S) \subseteq \bigcup_{d \in D} \overline{T_S^{\mathbb{R}[\underline{X}]_d}} \subseteq \overline{T_S^E}.$$

The following is a counterpart to Corollary 1.

Corollary 2. $\sum \mathbb{R}(\underline{X})^2$ is closed in $\mathbb{R}(\underline{X})$ with respect to \mathcal{T}_{fin} .

Proof. Every globally nonnegative rational function is a sum of squares of rational functions. \square

Example. In $\mathbb{R}(\underline{X}; \emptyset)$, the sums of squares are not closed. It is a long known fact that there are nonnegative polynomials which can not be written as a sum of squares of rational functions without poles. See for example [20, Lemma 1.1] or [18, Paragraph 6]. But as we have seen, these polynomials belong to the closure of the sums of squares in $\mathbb{R}(\underline{X}; \emptyset)$.

6. EXTENSION TO FINITELY GENERATED \mathbb{R} -ALGEBRAS

The aim of this section is to extend Theorems 1, 2 and 4 from polynomial rings to finitely generated \mathbb{R} -algebras.

Let A be a finitely generated \mathbb{R} algebra. A *character* of A is a unital \mathbb{R} -algebra homomorphism from A to \mathbb{R} . Write $V(A)$ for the set of all characters of A . Elements from A define functions on $V(A)$ by $a(\chi) := \chi(a)$. We equip $V(A)$ with the coarsest topology making all these functions continuous. One can embed $V(A)$ into some \mathbb{R}^n , by choosing generators x_1, \dots, x_n of A and sending $\chi \mapsto (\chi(x_1), \dots, \chi(x_n))$. In that case, the topology on $V(A)$ coincides with the usual topology from \mathbb{R}^n (see for example [11, Section 5.7]).

Let D be a multiplicative subset of $A \setminus \{0\}$ which contains 1. There is a canonical homomorphism $\iota: A \rightarrow F = D^{-1}A$, which however is not necessarily one-to-one. For every $d \in A$ write $\mathcal{Z}_A(d) = \{\chi \in V(A) \mid \chi(d) = 0\}$. Note that the set $V(A) \setminus \bigcup_{d \in D} \mathcal{Z}_A(d)$ consists of all characters of A which can be extended (via ι) to a character of F .

Let S be a finite subset of A . Write $K_S^A = \{\chi \in V(A) \mid \chi(S) \geq 0\}$. Let T_S^F be the preordering in F generated by the set $\iota(S)$.

For every subset \mathcal{X} of $V(A)$ write $\text{Pos}(\mathcal{X})$ for the set of all $a \in A$ such that $\chi(a) \geq 0$ for every $\chi \in \mathcal{X}$ and write $\text{Pos}^F(\mathcal{X})$ for the set of all elements $R \in F$ which have a representation $R = \frac{a}{d}$ with $a \in A$, $d \in D$ and $ad \in \text{Pos}(\mathcal{X})$.

Theorem 5. *Assume that A, D, S, F are as above. Also assume that there is an element $p \in D$ such that $p-1 \in \text{Pos}(K_S^A)$ and $kp - \sum_{i=1}^n x_i^2 \in \text{Pos}(K_S^A)$ for some integer $k \geq 1$ and generators x_1, \dots, x_n of A . Then*

- (1) *The closure of T_S^F in (F, \mathcal{T}_ω) is $\text{Pos}^F(K_S^A \setminus \bigcup_{d \in D} \mathcal{Z}_A(d))$.*
- (2) *The closure of T_S^F in $(F, \mathcal{T}_{\text{fin}})$ is $\text{Pos}^F(K_S^A)$.*
- (3) *For every linear functional L on F such that $L(T_S^F) \geq 0$, there exists a measure μ on $\overline{K_S^A \setminus \bigcup_{d \in D} \mathcal{Z}_A(d)}$ such that*

$$L\left(\frac{f}{d}\right) = \int \frac{f}{d} d\mu \quad \text{for every } \frac{f}{d} \in F.$$

Note that the set $\text{Pos}^F(K_S^A \setminus \bigcup_{d \in D} \mathcal{Z}_A(d))$ consists of all $R \in F$ such that $\tau(R) \geq 0$ for every character τ of F which satisfies $\tau(\iota(S)) \geq 0$.

Proof. Let $\pi: \mathbb{R}[\underline{X}] \rightarrow A$ be the unital \mathbb{R} -algebra homomorphism defined by $X_i \rightarrow x_i$ for $i = 1, \dots, n$ and $\pi^*: V(A) \rightarrow \mathbb{R}^n$ the corresponding embedding defined by $\chi \mapsto (\chi(x_1), \dots, \chi(x_n))$. If f_1, \dots, f_t are generators of the ideal $I = \ker \pi$, then $\pi^*(V(A)) = K_{\{\pm f_1, \dots, \pm f_t\}}$. Similarly, for a given set $S = \{g_1, \dots, g_m\} \subseteq A$ take elements $\tilde{g}_i \in \mathbb{R}[\underline{X}]$ such that $\pi(\tilde{g}_i) = g_i$ and note that

$$\pi^*(K_S^A) = K_{\tilde{S}} \quad \text{where} \quad \tilde{S} := \{\tilde{g}_1, \dots, \tilde{g}_m, \pm \tilde{f}_1, \dots, \pm \tilde{f}_t\}.$$

Let $\tilde{D} := \pi^{-1}(D)$ denote the localization $\tilde{D}^{-1}\mathbb{R}[\underline{X}]$ by E . π extends uniquely to a homomorphism $\tilde{\pi}: E \rightarrow F$, making the following diagram commutative:

$$\begin{array}{ccc} E & \xrightarrow{\tilde{\pi}} & F \\ \uparrow & & \uparrow \iota \\ \mathbb{R}[\underline{X}] & \xrightarrow{\pi} & A \end{array}$$

The preorderings $T_{\tilde{S}} \subseteq \mathbb{R}[\underline{X}]$ and $T_{\tilde{S}}^E \subseteq E$ do not depend on the specific choice of the \tilde{f}_i, \tilde{g}_i but only on I and S . We have

$$(*) \quad \tilde{\pi}(T_{\tilde{S}}^E) = T_S^F.$$

Now suppose D contains an element p with the properties required in Theorem 5. Any preimage \tilde{p} of p under π will have the corresponding property with respect to $K_{\tilde{S}}$.

For any linear functional L on F with $L(T_S^F) \geq 0$, the functional $\tilde{L} := L \circ \tilde{\pi}$ fulfills $\tilde{L}(T_S^E) \geq 0$, and is therefore integration on $\overline{K_{\tilde{S}} \setminus \bigcup_{\tilde{d} \in \tilde{D}} \mathcal{Z}(\tilde{d})}$ by Theorem 2. It follows that L is an integration on

$$\overline{K_S^A \setminus \bigcup_{d \in D} \mathcal{Z}(d)} = (\pi^*)^{-1} \left(\overline{K_{\tilde{S}} \setminus \bigcup_{\tilde{d} \in \tilde{D}} \mathcal{Z}(\tilde{d})} \right).$$

So we have proved assertion (3) of Theorem 5. Assertions (1) and (2) of Theorem 5 follow from Theorems 1 and 4, observations

$$\begin{aligned} \tilde{\pi}(\text{Pos}^E(K_{\tilde{S}} \setminus \bigcup_{\tilde{d} \in \tilde{D}} \mathcal{Z}(\tilde{d}))) &= \text{Pos}^F(K_S^A \setminus \bigcup_{d \in D} \mathcal{Z}_A(d)), \\ \tilde{\pi}(\text{Pos}^E(K_{\tilde{S}})) &= \text{Pos}^F(K_S^A) \end{aligned}$$

and (*) and from the following Lemma. □

Lemma 14. *Let E and F be \mathbb{R} -vector spaces, $\tilde{\pi}: E \rightarrow F$ a linear mapping which is onto and C a convex cone in E which contains $\ker \tilde{\pi}$. Then*

$$\tilde{\pi}(\overline{C}) = \overline{\tilde{\pi}(C)},$$

where either E, F are both equipped with \mathcal{T}_ω or both with \mathcal{T}_{fin} .

Proof. Suppose that E, F are equipped with \mathcal{T}_ω . The inclusion $\tilde{\pi}(\overline{C}) \subseteq \overline{\tilde{\pi}(C)}$ follows from the fact that $\tilde{\pi}$ is continuous in \mathcal{T}_ω . To prove the opposite inclusion, pick $\tilde{\pi}(e) \in \overline{\tilde{\pi}(C)}$. Then $L(\tilde{\pi}(e)) \geq 0$ for every linear functional L on F such that $L(\tilde{\pi}(C)) \geq 0$. Every linear functional L' on E such that $L'(C) \geq 0$ factors through $\tilde{\pi}$ because $\ker \tilde{\pi} \subseteq C$. It follows that $L'(e) \geq 0$ for every linear functional L' on E such that $L'(C) \geq 0$. So, $e \in \overline{C}$ which implies that $\tilde{\pi}(e) \in \overline{\tilde{\pi}(C)}$.

Suppose now that E, F are equipped with \mathcal{T}_{fin} . First note that by the same argument as in Lemma 9, the closure of a convex cone with respect to \mathcal{T}_{fin} is again a convex cone. Now the formula $\tilde{\pi}(\overline{C}) \subseteq \overline{\tilde{\pi}(C)}$ follows from the fact that $\tilde{\pi}$ is continuous. To prove the opposite inclusion it suffices to show that $\tilde{\pi}(C)$ is closed for every closed cone C containing $\ker \tilde{\pi}$. Suppose that C is such a cone and pick a finite-dimensional subspace W of F . Let V be a finite-dimensional subspace of E such that $\tilde{\pi}(V) = W$. By assumption, $C \cap V$ is closed in V in the Euclidean topology, hence also in the finest locally convex topology. Since $C \cap V$ is a closed cone in V which contains $V \cap \ker \tilde{\pi} = \ker \tilde{\pi}|_V$, it follows by the first paragraph (applied to $\tilde{\pi}|_V: V \rightarrow W$ instead of $\tilde{\pi}: E \rightarrow F$) that $\tilde{\pi}(C \cap V)$ is closed in $W = \tilde{\pi}(V)$. It remains to show that $\tilde{\pi}(C \cap V) = \tilde{\pi}(C) \cap \tilde{\pi}(V)$. Pick $\tilde{\pi}(e) \in \tilde{\pi}(C) \cap \tilde{\pi}(V)$. We have that $e = c + i = v + j$ for some $c \in C$, $v \in V$ and $i, j \in \ker \tilde{\pi}$. Since $\ker \tilde{\pi} \subseteq C$, $c + i - j = v$

belongs to $C \cap V$. It follows that $\tilde{\pi}(e) = \tilde{\pi}(v)$ belongs to $\tilde{\pi}(C \cap V)$. The opposite inclusion is clear. \square

It would be interesting to know whether Theorem 5 also holds for the algebra $A = \mathcal{O}(\mathbb{R}^n)$ of analytic functions on \mathbb{R}^n . In this case we can use [1, Theorem 2.4] instead of [22, Theorem 5.1], so the assumption $p \in D$ for certain p may not be necessary.

REFERENCES

- [1] F. Acquistapace, C. Andradas, F. Broglia, *The strict Positivstellensatz for global analytic functions and the moment problem for semianalytic sets*, Math. Ann. **316** (2000), pp. 609-616. MR1758445 (2001g:14087)
- [2] N. I. Akhiezer, *The Classical Moment Problem and Some Related Questions in Analysis*, Oliver & Boyd, Edinburgh/London, 1965. MR0184042 (32 #1518)
- [3] T. M. Bisgaard, *The topology of finitely open sets is not a vector space topology*, Arch. Math. **60** (1993), pp. 546-552. MR1216700 (94g:46013)
- [4] J. Bochnak, M. Coste, M.-F. Roy, *Real Algebraic Geometry*, Erg. Math. Grenzgeb. 36, Springer, Berlin (1998). MR1659509 (2000a:14067)
- [5] N. Bourbaki, *Topological vector spaces, Chapters 1-5*, English edition, Springer Verlag, Masson 1987. MR0910295 (88g:46002)
- [6] A. Bultheel, P. González-Vera, E. Hendriksen, O. Njåstad, *Orthogonal rational functions*, Cambridge Monographs on Applied and Computational Mathematics vol. 5, Cambridge University Press, Cambridge, 1999. MR1676258 (2000c:33001)
- [7] A. Bultheel, P. González-Vera, E. Hendriksen, O. Njåstad, *Elements of a theory of orthogonal rational functions*, Rev. Acad. Canaria Cienc. **11** (1999), pp. 127-152. MR1780786 (2001i:42041)
- [8] J. D. Chandler, Jr., *Rational moment problems for compact sets*, J. Approximation Theory **79** (1994), pp. 72-88. MR1294321 (95h:44019)
- [9] S. Kuhlmann, M. Marshall, *Positivity, sums of squares and the multidimensional moment problem*, Trans. Amer. Math. Soc. **354** (2002), pp. 4285-4301. MR1926876 (2003j:14078)
- [10] S. G. Krantz, H. R. Parks, *Geometric Integration Theory*, <http://www.math.wustl.edu/~sk/books/root.pdf>.
- [11] M. Marshall, *Positive polynomials and sums of squares*, AMS Surveys and Monographs vol. 146, AMS, Providence, 2008. MR2383959
- [12] M. Marshall, *Extending the Archimedean Positivstellensatz to the non-compact case*, Canad. Math. Bull. **44** (2001), pp. 223-230. MR1827856 (2002b:14073)
- [13] M. Marshall, *Approximating positive polynomials using sums of squares*, Canad. Math. Bull., **46** (2003), pp. 400-418. MR1994866 (2004f:14084)
- [14] T. W. Palmer, *Banach algebras and the general theory of *-algebras, Vol. 2*, Encyclopedia of Mathematics and its Applications, 79. Cambridge University Press, Cambridge, 2001. pp. i-xii and 795-1617. MR1819503 (2002e:46002)
- [15] A. Prestel, C. N. Delzell, *Positive polynomials*, Springer, Berlin (2001). MR1829790 (2002k:13044)

- [16] M. Putinar, F. -H. Vasilescu, *Solving moment problems by dimensional extension*, Ann. of Math. (2) **149** (1999), pp. 1087-1107. MR1709313 (2001c:47023b)
- [17] B. Reznick, *Uniform denominators in Hilbert's Seventeenth Problem*, Math. Z. **220** (1995), pp. 75-97. MR1347159 (96e:11056)
- [18] B. Reznick, *Some concrete aspects of Hilbert's 17th problem*, in: Real Algebraic Geometry and Ordered Structures, (C. N. Delzell, J.J. Madden eds.) Cont. Math. vol. 253 (2000), pp. 251-272. MR1747589 (2001i:11042)
- [19] W. Rudin, *Real and Complex Analysis*, 3rd ed., McGraw-Hill 1987. MR0924157 (88k:00002)
- [20] C. Scheiderer, *Sums of squares of regular functions on real algebraic varieties*, Trans. Am. Math. Soc. **352** (1999), pp. 1039-1069. MR1675230 (2000j:14090)
- [21] K. Schmüdgen, *The K -moment problem for compact semi-algebraic sets*, Math. Ann. **289** (1991), pp. 203-206. MR1092173 (92b:44011)
- [22] M. Schweighofer, *Iterated Rings of Bounded Elements and Generalizations of Schmüdgen's Positivstellensatz*, J. Reine Angew. Math. **554** (2003), pp. 19-45. MR1952167 (2004b:13028) (unpublished Erratum available at <http://perso.univ-rennes1.fr/markus.schweighofer/publications/irobeerr.pdf>)
- [23] J. Stochel, F. H. Szafraniec, *Algebraic operators and moments on algebraic sets*, Portugal. Math. **51** (1994), 25-45. MR1281954 (95d:47013)
- [24] F. -H. Vasilescu, *Spaces of fractions and positive functionals*, Math. Scand. **96** (2005), 257-279. MR2153414 (2007b:47040)
- [25] J. H. Williamson, *On Topologizing the field $\mathbb{C}(t)$* , Proc. Amer. Math. Soc. **5** (1954), pp. 729-734. MR0063574 (16,145f)

CIMPRIČ, JAKA, FACULTY OF MATHEMATICS AND PHYSICS,
 UNIVERSITY OF LJUBLJANA, JADRANSKA 21, SI-1000 LJUBLJANA, SLOVENIJA,
 EMAIL: CIMPRIC@FMF.UNI-LJ.SI

MARSHALL, MURRAY, DEPARTMENT OF MATHEMATICS AND STATISTICS,
 UNIVERSITY OF SASKATCHEWAN, SASKATOON, SK S7N 5E6, CANADA,
 EMAIL: MARSHALL@MATH.USASK.CA

NETZER, TIM, FACHBEREICH MATHEMATIK UND INFORMATIK,
 UNIVERSITÄT LEIPZIG, D- 04009 LEIPZIG, GERMANY,
 E-MAIL TIM.NETZER@MATH.UNI-LEIPZIG.DE