

**Special Session on Real
Algebraic Geometry and
Applications.
MTNS 2010, Budapest,
Hungary, July 5–9 2010.**

July 8, 2010

Salma Kuhlmann
Schwerpunkt Reelle Geometrie und Algebra,
Fachbereich Mathematik und Statistik,
Universität Konstanz,
78457 Konstanz, Germany

Email: salma.kuhlmann@uni-konstanz.de

The slides of this talk are available at:

<http://math.usask.ca/~skuhlman/slidesmtns2010.pdf>

Positivity in Power Series Rings.

Preliminaries.

- Let $\mathbb{R}[X] := \mathbb{R}[X_1, \dots, X_n]$ be the ring of polynomials in n variables and real coefficients.
- A subset $M \subseteq \mathbb{R}[X]$ is a **quadratic module** if $1 \in M$, M is closed under addition and multiplication by squares (i.e. $a^2 f \in M$, $\forall a \in \mathbb{R}[X]$ and $f \in M$).
- A **quadratic preordering** is a quadratic module which is also closed under multiplication.
- The smallest preordering of $\mathbb{R}[X]$ is the set of **sums of squares** of $\mathbb{R}[X]$, denoted by $\Sigma \mathbb{R}[X]^2$.

- Given a finite subset $S = \{f_1, \dots, f_s\}$ of $\mathbb{R}[X]$, the smallest preordering of $\mathbb{R}[X]$ containing S (**preordering finitely generated by S**) is:

$$T_S = \left\{ \sum_{e \in \{0,1\}^s} \sigma_e f^e : \sigma_e \in \sum \mathbb{R}[X]^2, f_1, \dots, f_s \in S \right\}$$

where $f^e := f_1^{e_1} \cdots f_s^{e_s}$, if $e = (e_1, \dots, e_s)$.

- Let $S = \{f_1, \dots, f_s\} \subset \mathbb{R}[X]$, S defines a **basic closed semialgebraic** subset of \mathbb{R}^n :

$$K = K_S = \{x \in \mathbb{R}^n : f_1(x) \geq 0, \dots, f_s(x) \geq 0\}$$

- Consider polynomials **positive semi-definite** on K_S :

$$\text{Psd}(K_S) := \{f \in \mathbb{R}[X] : f(x) \geq 0 \text{ for all } x \in K_S\}$$

- $\text{Psd}(K_S)$ is a preordering in $\mathbb{R}[X]$ and $T_S \subseteq \text{Psd}(K_S)$.

Hilbert's 17th Problem is concerned with the issue of representation of positive semi-definite polynomials; motivated by the question: when is it true that $\text{Psd}(K_S) = T_S$?

We say:

- T_S is **saturated** if $\text{Psd}(K_S) = T_S$.

Low dimensional compact sets

In [S1] Scheiderer showed:

Theorem 0.1 *If $\dim(K_S) \geq 3$, then there exists a polynomial $p(X) \in \mathbb{R}[X]$ such that $p(x) \geq 0$ for all $x \in \mathbb{R}^n$ but $p \notin T_S$ (so T_S cannot be saturated).*

In the same paper, he also shows:

Theorem 0.2 *If $n = 2$ and K_S contains a cone of dimension 2, then there exists a polynomial $p(X) \in \mathbb{R}[X]$ such that $p(x) \geq 0$ for all $x \in \mathbb{R}^n$ but $p \notin T_S$ (so T_S cannot be saturated).*

This left open the question formulated in Kuhlmann – Marshall [K-M]: Are there compact examples of $K_S \subseteq \mathbb{R}^2$ for which T_S is saturated?

Schmüdgen's Positivstellensatz and Scheiderer's Local Global Principle

In [Sc1] Schmüdgen proved the following Striktpositivstellensatz:

Theorem 0.3 *If K_S is compact, then $f > 0$ on K_S implies that $f \in T_S$.*

Note: the hypothesis $f > 0$ cannot be replaced by $f \geq 0$.

Recently, Scheiderer developed in a series of papers [S2], [S3], [S4] several *local global principles* to determine when a polynomial $f \geq 0$ on K_S belongs to T_S . His results generalize Schmüdgen's Striktpositivstellensatz.

Scheiderer's LGP states:

Theorem 0.4 *Suppose $f, g_1, \dots, g_s \in \mathbb{R}[X]$, assume that K_S is compact, $f \geq 0$ on K , and f has just finitely many zeros in K . Then the following are equivalent:*

1. $f \in T_S$.
2. *For each zero p of f in K , f lies in the preordering of the completion of $\mathbb{R}[X]$ at p generated by g_1, \dots, g_s .*

Where the **completion** of $\mathbb{R}[X]$ at $p = (p_1, \dots, p_n) \in \mathbb{R}^n$ is the ring of formal power series $\mathbb{R}[[X_1 - p_1, \dots, X_n - p_n]]$.

Saturation for compact affine two dimensional sets.

In the two dimensional affine compact case his LGP allows him to show that certain finitely generated preorderings are saturated We need the following definitions to state the theorem. We say that p is a **singular** zero of g if

$$\delta g/\delta X(p) = \delta g/\delta Y(p) = 0 .$$

We say that g and h **meet transversally** at p if p is a non-singular zero of both g and h , and the curves $g = 0$ and $h = 0$ have distinct tangents at p .

Theorem 0.5 *Let $S = \{g_1, \dots, g_s\}$ be irreducible polynomials in $\mathbb{R}[X, Y]$. Suppose that $K = K_S \subseteq \mathbb{R}^2$ is compact, and, for each boundary point p of K , either*

- 1. there exists i such that p is a non-singular zero of g_i , and K is defined locally at p by the single inequality $g_i \geq 0$; or*
- 2. there exists i, j such that p is a non-singular zero of g_i and g_j , g_i and g_j meet transversally at p , and K is defined locally at p by $g_i \geq 0, g_j \geq 0$.*

Then the preordering of $\mathbb{R}[X, Y]$ generated by g_1, \dots, g_s is saturated.

With these tools, he was able to produce examples:

Example 0.6

- The preordering generated by:
 $S = \{1 + X, 1 - X, 1 + Y, 1 - Y\}$ (K_S is the unit square)
is saturated.
- The preordering generated by:
 $S = \{1 - X^2 - Y^2\}$ (K_S is the unit disk) is saturated.
- $S = \{X^3 - Y^2, 1 - X\}$ then $X \geq 0$ on K_S but $X \notin T_S$
($(0, 0)$ is a singular zero of $X^3 - Y^2$).
- $S = \{Y, Y - X^3, 1 - X\}$, $X \geq 0$ on K_S but $X \notin T_S$
(the curves $Y = 0$ and $Y = X^3$ do not meet transversally
at $(0, 0)$).

In [C-K-M] we extend Scheiderer's theorem to handle certain cases where the boundary curves do not meet transversally, for example our theorem covers the following examples:

Example 0.7 $S = \{X, 1 - X, Y, X^2 - Y\}$ or $S = \{1 + X, 1 - X, Y, X^2 - Y\}$. In these examples, the boundary curves $Y = 0$ and $Y = X^2$ share a common tangent at the origin, so Scheiderer's result does not apply. The fact that saturation holds in these examples is a consequence of our main result, Corollary 0.8 in the next slide.

C–K–M Generalization of Scheiderer's Theorem.

Corollary 0.8 *Let $S = \{g_1, \dots, g_s\}$ be irreducible polynomials in $\mathbb{R}[X, Y]$. Suppose that $K = K_S \subseteq \mathbb{R}^2$ is compact, and, for each boundary point p of K , either*

- 1. there exists i such that p is a non-singular zero of g_i , and K is defined locally at p by the single inequality $g_i \geq 0$; or*
- 2. there exists i, j such that p is a non-singular zero of g_i and g_j , g_i and g_j meet transversally at p , and K is defined locally at p by $g_i \geq 0, g_j \geq 0$; or*

3. *there exists i, j such that p is a non-singular zero of g_i and g_j , g_i and g_j share a common tangent at p but do not cross each other at p , and K is described locally at p as the region between $g_i = 0$ and $g_j = 0$; or*
4. *there exists i, j, k such that p is a non-singular zero of g_i, g_j and g_k , g_i and g_j share a common tangent at p , g_i and g_k meet transversally at p , and K is described locally at p as the part of the region between $g_i = 0$ and $g_j = 0$ defined by $g_k \geq 0$.*

Then the preordering of $\mathbb{R}[X, Y]$ generated by g_1, \dots, g_s is saturated.

The End

References

- [C-K-M] J. Cimpric, S. Kuhlmann and M. Marshall: *Positivity in Power Series Rings*, Advances in Geometry **10** (2010).
- [K-M] S. Kuhlmann and M. Marshall: *Positivity, sums of squares and the multi-dimensional moment Problem*, TAMS **354** (2002), 4285–4301
- [S1] C. Scheiderer: *Sums of squares of regular functions on real algebraic varieties*, Trans. Amer. Math. Soc. **352**, 1030–1069 (1999)
- [S2] C. Scheiderer: *Sums of squares on real algebraic curves*, Math. Zeit. **245** (2003), 725–760
- [S3] C. Scheiderer: *Distinguished representations of non-negative polynomials*, J. Algebra **289** (2005), 558–573
- [S4] C. Scheiderer: *Sums of squares on real algebraic surfaces*, Manuscripta Math. **119** (2006), 395–410
- [Sc1] K. Schmüdgen: *The K -moment problem for compact semi-algebraic sets*, Math. Ann. **289**, 203–206 (1991)